

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification :  Not classified		A2	(11) International Publication Number: <b>WO 00/41524</b>
			(43) International Publication Date: 20 July 2000 (20.07.00)
(21) International Application Number: PCT/US00/00580 (22) International Filing Date: 10 January 2000 (10.01.00) (30) Priority Data: 60/115,498 11 January 1999 (11.01.99) US (71) Applicant (for all designated States except US): PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TABOR, Stanley [US/US]; 9a Lowell Street, Cambridge, MA 02138 (US). RICHARDSON, Charles [US/US]; 78 Chestnut Hill Road, Chestnut Hill Road, MA 02167 (US). (74) Agent: LYON & LYON LLP; 633 West Fifth Street, Suite 4700, Los Angeles, CA 90071-2066 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published Without international search report and to be republished upon receipt of that report.	
(54) Title: ISOTHERMAL AMPLIFICATION OF DNA			
(57) Abstract			
<p>A method of amplifying a template DNA molecule comprising incubating the template DNA molecule in a reaction mixture comprising a DNA polymerase and at least one accessory protein at a constant temperature to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers and the template DNA molecule does not have terminal protein covalently bound to either 5' end.</p>			
<pre> 10      20      30      40      50      60      70      80 ATGACGATT  GCGACGATTC  CGATGATGTA  TTCTTTTACC  ACATTCCTTG  TGACAACTGT  GGGAGTAGTG  ATGGGAATCT  80 GCTGTCTCT  GACGGACACA  CGTCTCTCTA  CGATATCCAG  AGGTGGACTG  CTGTATATGA  AGACACTAAA  GAGAGGCTTT  160 CAAACTCGAA  ACCCTCCTGG  GGAAGGCTCG  GGACTTACAA  COTGTGGAAC  TTGGGGGAAT  CCAATGGACG  CTACTCCGCG  240 TTAAGTCGGA  GAGGAAATCT  CAAAGAAACC  TGTGAGAAGG  CTGGCTACTG  GATTGCCAAA  GTAGACGGTG  TGATGTACCA  320 AGTGGCTGAC  TATCGGGACC  AGAACGGCAA  CATTGTGAGT  CAGAAGGTTG  GAGATAAGA  TAGAACTTTC  AGACCACTG  400 GTATGCACA  GACTGAGCTG  CTGTTCGGGA  AGCACTTGTG  GAATGGGGTG  AGAAGATTG  TGTATACAGA  AGGTGAATTC  480 GACATGCTTA  CGGTGATGGA  ACTTCAAGAC  TGTAAATATC  CTGTAGTGTG  GTTGGGTGAC  GGTGCTCTGT  CGGTGAGAA  560 GACATGCTGT  GCGAACTAGC  AATACTTTGA  CCAATTCGAA  CAGATTATCT  TAATGTTTGA  TATGAGCAAA  GCAGGGCGCA  640 AAGCAATGGA  AAGAGCTGCA  CAGGTTCTAC  CTGCTCGTAA  GGTACGAGTG  GCAATCTCTG  CGTGTAAAGA  TGCAAAACAG  720 TGTCACTTAA  ATGCTCACGA  CGGTGAATTC  ATGGAGCAAG  TGTGAATGC  TGTCTCTTGG  ATTCTGTATG  GTGTGTATTC  800 GGCTCTTTCG  TTACGTGAA  GAATCGCTGA  GCACCTATCG  TCGAGGAAT  CAGTAAGTTT  ACTTTCTAGT  GGTCTCACTG  880 GTATCAGCA  TANGACCTTA  GGTGCGCTG  GTGTGAAAT  CATTATGCT  ACTTCCGTT  CCGGTATGCG  TAAATCAAGC  960 TTCTGCTGCT  AACAGCTCT  ACAAATGGGC  AAGCGATGCG  GCAAGAGGT  AGCTTACGCG  ATGCTTAGCG  AGTCCCTTGA  1040 GGAGACCGCT  GAGGACCTTA  TAGGTCTACA  CAACCGTGT  CGACTGAGAC  AATCCGACT  ACTAAACAGA  GAGATTATG  1120 AGAACGGTAA  GTTGCACCA  TGGTTCGATG  AACTGTTGCG  CAACGATCG  TTGCTCTTAT  ATGACTCTAT  CGCGAAGGCT  1200 GAGACGGATA  GACTGCTCG  TAAGTGTGCG  TACATGCGCT  CAGGCTTGG  CTGTGACGTA  ATGACTCTAG  ACCCAATCTC  1280 AATCTGCTTA  TCCGCTCTGT  GTGAATCCGA  TGAGCTAAG  ATGATTAACA  ACTGATGAC  CAGCTCTGAA  GGGTCCCTTA  1360 AGTCACTGCG  GTGTGCTGT  GTGGAATTT  GTCACTTAA  GAACCCAGAC  AAGGTAAAG  CACATAGCA  AGGTCCCGCC  1440 GTCTCTATTA  CTGACTTACG  TGGTCTGCG  GCACTACGCC  AATCTATGTA  TACTATATT  GCGCTTAGCG  GTATCTACCA  1520 AGCGATATG  CTTAAGCTG  TCGTGTGCG  TATCTCTAG  TGCGCTTTA  CTGTGTATAC  TGGTATCGCT  GGTCACTAGT  1600 ATACACAA  GGAACCGGA  TGCTGTGAC  CATCAAGTA  CTGAGGGAA  GAGAGTCA  ACTCAGATG  AACAGCTGCG  1680 TCCACGACA  CTGACTCTGT  ACAGGATCT  TGATGACTT  CCGACGACT  ACGAGAGTT  TCGCTCGGA  GTCTCATCTT  1760 AATGAGCTC  ACTAAGGGA  ACACACATG  TTCAACTGA  TTAAGAGTT  AGGCCAATC  CTGCTCTGTA  TGTACACGTT  1840 GGAAGCCAG  CGACTGAAG  ATGAGGCTG  TAAAGAGGCC  ACACGATCA  GCGCTCTGCG  GATGTGCTC  AAACCTGTT  1920 TTGCGCTTAC  CCGACACAC  AGCGGATTC  CGCTTTTCCA  TTGAGCTGT  TTCTCTGCG  GACCTCTGCG  GCGGCTGTGT  2000 TGTGATCCA  TGTGATTTCT  CTGTGAGTT  AGCTTTGCTG  GTGTGTGGA  GTTCTAGTCC  TGACGAGAA  CCGCGCGGGA  2080 TTGAGCAAT  GCGAGCTAAT  CCGGAATCG  ACTTACGCG  AATGCTTCT  TTCTATAC  AGACCCGAAA  GCTCTCTGCT  2160 TTGATGCTG  GCTCTTCTA  GCGCTTAAT  TTAAGAGCG  TCACCTTCA  GTGTGCTAGT  GCGTCTGCT  GATGTGCTCA  2240 GTATCAGCG  CAGTGTGAT  TATGTACCA  CCGCGAGGA  TAATTTATCA  CCGCAGATGG  TTATCTGTAT  GTTTTGTATA  2320 TGAATTTAT  TTTTGCAGG  GCGCATGTT  TGTGATGTA  GAGATCGCG  TGTATACAA  CCGCGAAGG  AAGCTGAGTT  2400 GCTGTGCTG  ACCTGTGAG  AATACTAGC  ATAACCCCT  ACCGGCTTA  ACGAGCTAT  GCTCTAGCA  TCCAGGGTGA  2480 GAGAACATAT  ATCCGATAT  CCGCAAGAG  CCGCGCGAG  ACAGGTGCG  TGACTGCTT  AGCAATTTAA  TCACTATAGG  2560 CGACCGATT  AAGCTTGGC  CGCGACATG  ACGAACCTT  CGATTCGTA  TCCCGGAAA  TTAATACGAG  TCACTATAGG  2640 GAGAACCTTA  AAGGCTCTT  AGGGGTTTT  TTGCTGAAG  CTTTAAAGAG  GAGATATACA  TATGCTGATA  GCAATCCGCT  2720 AGCACCTAT  GTCCGAGAA  TCAATATGCT  ATAGACCTT  TGGCTGACT  GGTATCAAG  ATAGACCTT  AGCTGCGGCT  2800 GGTGTGAGG  TCATTATGCT  CACTTCCGT  TCCGATATG  GTAAGTCAAC  GTTCTGCT  CAACACCTG  TACATAGGGG  2880 CAGACCGAT  CGAGAGAGG  TAGGCTAGC  GATGTGAG  GAGTCCGTT  AGGAGACCG  TGAAGACCT  ATAGGTCTAC  2960 ACAACTGAT  CGACTGAAG  CAATCCGACT  CACTAAGAG  AGAGATTAT  GAGAACGGA  AGTTGAGCA  ATGTTCTGAT  3040 GACTGTTCG  CGAACGATAC  GTTCACTCT  TATGACTAT  TCGCCGAGC  TGACAGGAT  AGATCTGCG  CTAACTGCG  3120 CTACATGCG  TCGCGTTGG  GTGTGACGT  AATCACTTA  GACCAATCT  CAATCTGCT  ATCCGCTCT  GGTGAATCCG  3200 ATAGCGTAA  GATGATGAG  AACTGATGA  CCAAGCTCA  AGGTTGCT  AGTCAACTG  GTGTGCTCT  GGTGTAAAT  3280 TGTCACTTA  AGAACCCGA  CAAAGTAAA  GCACATAGG  AAGTCCGCT  CGTTCTAT  ACTGACCTAC  GTGTGCTG  3360 CGCACTACG  CAATCTATG  ATACTATTAT  TGCCCTTGA  CTAATACG  AAGCGATAT  GCTCACTT  CTCTCTGCT  3440 </pre>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 00/41524

PCT/US00/00580

## ISOTHERMAL AMPLIFICATION OF DNA

### FIELD OF THE INVENTION

The present invention relates to the field of amplification of DNA and RNA.

### BACKGROUND OF THE INVENTION

5           This invention was made with government support including a grant from the U.S. Dept. of Energy, contract number DE-FG02-96ER62251. The U.S. government may have certain rights in the invention.

The following is a discussion of the relevant art, none of which is admitted to be prior art to the appended claims.

10           A variety of methods have been developed to efficiently amplify DNA using purified proteins. These generally can be classified as ones that use thermocycling of the reaction mixtures and ones in which the reactions are isothermal. Methods that use thermocycling of the reaction mixture are based on the polymerase chain reaction, or PCR (Saiki *et al.*, Science 230:1350, 1985). In PCR, two primers are added to the target  
15   DNA, such that the two primers are complementary to opposite strands of the target sequence with their 3' ends oriented towards one another. Multiple cycles of denaturation of target DNA, annealing of the two primers, and then extension of the primers with a DNA polymerase, results in an exponential synthesis of the target DNA sequence located between the two primers. This procedure requires the use of a DNA  
20   polymerase that is thermostable in order to survive the high temperatures required to denature the product DNA each cycle. The most common polymerase used for PCR is from *Thermus aquaticus*, or Taq DNA polymerase (Saiki *et al.*, Science 239:487, 1988). Modification of the reaction conditions and the enzyme mixture used can allow the amplification of DNA up to 50 kbp in length (Barnes, Proc. Natl. Acad. Sci. USA  
25   91:2216, 1994 and Cheng *et al.*, Proc. Natl. Acad. Sci. USA 91:5695, 1994).

WO 00/41524

2

PCT/US00/00580

Isothermal procedures for the amplification of nucleic acid include (1) Q $\beta$  replicase, (2) self-sustained sequence replication (3SR), (3) strand displacement amplification (SDA), (4) terminal protein-primed DNA amplification using  $\phi$ 29 DNA polymerase, and (5) rolling circle amplification (RCA).

- 5        The RNA-dependent RNA polymerase Q $\beta$  polymerase has been used for gene detection in a strategy that uses two probes that hybridize to adjacent positions on a target sequence that are then ligated to form an amplifiable reporter RNA (Tyagi *et al.*, Proc. Natl. Acad. Sci. USA 93:5395, 1996).

- 10       Self-sustained sequence replication (3SR) is a method of amplifying RNA that depends on the action of three enzymes; reverse transcriptase, DNA-dependent RNA polymerase and ribonuclease H (Guatelli *et al.*, Proc. Natl. Acad. Sci. USA 87:1874, 1990).

- 15       Strand displacement amplification (SDA) is a DNA amplification system that uses a restriction enzyme to introduce specific nicks in a target to be amplified and a DNA polymerase that is capable of strand displacement synthesis (Walker *et al.*, Proc. Natl. Acad. Sci. USA 89:392, 1992).

- 20       Terminal protein-primed DNA amplification exponentially amplifies linear  $\phi$ 29 DNA using the  $\phi$ 29 proteins DNA polymerase, terminal protein, double-stranded DNA binding protein, and single-stranded DNA binding protein (Blanco *et al.*, Proc. Natl. Acad. Sci. USA 91:12198, 1994). To date this method has only been successful using linear  $\phi$ 29 DNA as template.

- 25       Rolling circle amplification (RCA) can either be linear or exponential amplification of circular oligonucleotide probes that use two primers that anneal to each of the two strands, and a DNA polymerase that strand displaces (Lizardi *et al.*, Nature Genetics 19:225, 1998 and Zhange *et al.*, Gene 211:277, 1998).

In order to be used as a generalized amplification system, all of these methods require the addition of specific oligonucleotide primers that are extended by a

WO 00/41524

PCT/US00/00580

3

polymerase; the primers serve to fix the limits of the segment of nucleic acid to be amplified.

### SUMMARY OF THE INVENTION

The present invention provides an *in vitro* method to amplify DNA exponentially  
5 at a constant temperature and without the use of exogenously added primers by using a DNA polymerase and various accessory proteins. The accessory proteins function with the DNA polymerase to carry out amplification under these conditions.

In one aspect, the method involves the addition of a mixture of enzymes including a DNA polymerase, a helicase and a primase and single-stranded binding  
10 protein to template DNA, which can result in amplification of the template DNA of at least 10-fold and up to and greater than 1 million-fold amplification. Such an amplification system is useful for common molecular biology procedures, including but not limited to, the preparation of plasmid DNA from cells for DNA sequencing, cloning, providing substrates for transcription/translation reactions, and mutagenesis. In addition,  
15 this generalized system for DNA amplification can serve as a useful way to prepare large amounts of human genomic DNA from small samples for genotype analysis. This could be important both for preserving or "immortalizing" small amounts of rare DNA samples, and for more general use in clinical applications as a way to avoid removing large amounts of blood from patients to obtain DNA for genotyping. Finally, this  
20 amplification system provides a very sensitive detection system for DNA contamination in samples, for example in enzymes that have been purified under good manufacturing protocol (GMP) for use in human therapeutics.

One way in which the present invention is distinguished from other exponential procedures for amplification is that it does not require the use of exogenous  
25 oligonucleotide primers. Exogenous oligonucleotide primers refer to small nucleic acid molecules that are generally in the range of 6 to 50 bases in length, although longer primers can work. They are normally synthesized by chemical as opposed to enzymatic methods. The primers are complementary to certain regions of the template and are

WO 00/41524

PCT/US00/00580

4

utilized by the polymerase to initiate DNA synthesis. The region of the template that is amplified is defined by the primers utilized. Primers are required in the present invention, however they are made by one of the components of the system, the primase, *e.g.*, gene 4 protein, rather than being synthesized in advance and then added to the system. The addition of primers to the system of the present invention has little effect on increasing the rate of DNA synthesis (generally less than two-fold), as the system works very well even without exogenous primers. Thus, even if the replication system of the present invention contains exogenous primers, there is still substantial DNA synthesis that is exponential in nature that is independent of the primers. One of ordinary skill in the art would readily be able to distinguish DNA synthesis carried out by the claimed method from exogenously added primer dependent synthesis. In the situation where DNA synthesis is dependent on specific exogenously added primers, a new set of primers will be required for each new DNA template amplified. In the situation where DNA synthesis is dependent on the exogenous addition of random primers (*e. g.* random hexamers) the DNA synthesis will not be exponential. This invention describes a system whereby DNA synthesis is exponential even in the absence of any exogenously added primers.

The template DNA used in this invention also does not need have a terminal protein bound to either 5' end. A terminal protein is a protein like the type used in the bacteriophage  $\phi$  29 replication system, *e.g.*, phage  $\phi$  29 the gene 3 protein (Salas, Ann. Rev. Biochem. 60:39, 1991). Such a terminal protein may be present but is not functionally required for the amplification system to work.

The term "constant temperature" refers to an amplification reaction that is carried out under isothermal conditions as opposed to thermocycling conditions. Thermocycling conditions are used by polymerase chain reaction methods in order to denature the DNA and anneal new primers after each cycle. Constant temperature procedures rely on other methods to denature the DNA, such as the strand displacement ability of some polymerases or of DNA helicases that act as accessory proteins for some DNA polymerases. Thus, the term "constant temperature" does not mean that no temperature

WO 00/41524

5

PCT/US00/00580

fluctuation occurs, but rather indicates that the temperature variation during the amplification process is not sufficiently great to provide the predominant mechanism to denature product/template hybrids. Preferably the constant temperature is less than 60°C, more preferably less than 50°, still more preferably less than 45°, and can even be less than 40°C.

The constant temperature amplification is carried out "*in vitro*", meaning that the reaction is not carried out in cells. Generally the amplification is performed using purified or at least substantially purified proteins.

In a preferred embodiment the DNA polymerase is from a bacteriophage. Further preferred is that the DNA polymerase is from bacteriophage T7. An even further preferred embodiment is that the DNA polymerase is a mixture of enzymes including both the wild-type DNA polymerase from bacteriophage T7 as well as a mutant T7 DNA polymerase that lacks its 3' to 5' exonuclease activity. It is also preferred that the reaction mixture contains a helicase or a primase. The helicase and primase are preferably the gene 4 protein of bacteriophage T7, most preferably the 63-kDa T7 gene 4 protein that encodes the helicase/primase complex. Also preferred in the reaction mixture is a single-stranded DNA binding protein, preferably a single-stranded DNA binding protein from *Escherichia coli*.

Applicant has surprisingly found that such a combination of proteins results in an unexpectedly large amount of DNA synthesis, and that the kinetics of this DNA synthesis is exponential rather than linear in nature. By exponential it is meant that at some period of time during the reaction the rate of DNA synthesis increases. In other words, the amount of DNA synthesized at a particular time will be greater than twice the amount of DNA synthesized at half the time. For example, if the amount of DNA synthesized after 20 minutes is ten times the amount of DNA synthesized after 10 minutes, then the kinetics of DNA synthesis is exponential. On the other hand, if the amount of DNA synthesized after 20 minutes is only twice the amount of DNA synthesized after 10 minutes, then the kinetics of DNA synthesis is linear. In order to test for whether DNA synthesis is linear or exponential, a small amount of radioactive

WO 00/41524

6

PCT/US00/00580

deoxynucleoside triphosphate can be added to the mixture; *e.g.* example, one million cpm of [ $\alpha$ - $^{32}$ P]dATP in a 50  $\mu$ l reaction, or a specific activity of 40 cpm per pmol of dATP if the dATP concentration is 500  $\mu$ M. Once the reaction is initiated by the addition of the enzyme mixture, aliquots are removed at varying times and the amount of DNA synthesized is determined, as described in Example 1. For example, 5  $\mu$ l aliquots could be removed at 0, 1, 2, 4, 8, 16 and 32 min (although the amount of each aliquot removed and the times can vary). The amount of DNA synthesized at the different times is plotted as a function of the time of the reaction. If DNA synthesis is exponential, then the shape of the plot will be sigmoidal with the slope of the curve increasing over some portion of the time interval. A reaction will be exponential only for a portion of the time; the dNTPs will be rapidly used and the rate of new DNA synthesis will then rapidly decrease. Thus to determine if a reaction is exponential it is important to inspect the kinetics of DNA synthesis over a broad range of time (for example, from 0 to 30 min), and determine if the rate of DNA synthesis increases over any period of the time of the reaction. Also, the total amount of DNA synthesized is quite large, typically at least 10-fold more than the amount of template added to the reaction mixture, and preferably is at least 100-fold greater, or at least 1000-fold greater, or at least 10,000-fold greater, or at least 100,000-fold greater, or at least 1,000,000-fold greater, or at least 10,000,000-fold greater or even more. Thus, one unique feature of the present invention is that synthesis is both exponential and results in the production of a large amount of DNA, *e.g.* 10-100 times or more greater than the amount of template. Furthermore as the present invention does not require the use of exogenously added primers specific for each template being amplified, it provides a generalized amplification method not limited to the region of the DNA template delineated by primers.

Another very effective method of monitoring the amount of DNA synthesized using the isothermal exponential amplification system is to use fluorescent probes or fluorescent dyes such as SYBR Green II (Molecular Probes, Eugene, OR) to continuously report the amount of DNA in the reaction in real time. The basic principle of this method has been used successfully to determine the initial concentrations of specific RNAs and DNAs in a PCR reaction (Gibson *et al.*, Genome Research 6:995, 1996). Fluorescence can be monitored at intervals of 15 seconds using fluorescence-



WO 00/41524

PCT/US00/00580

7

based real-time PCR instruments such as the PE Biosystems 5700 (PE Biosystems, Foster City, CA), the Roche LightCycler (Indianapolis, IN), or fluorescence microtiter plate readers that can maintain a constant temperature and can carry out kinetic measurements such as the SPECTRAmax (Molecular Devices, Sunnyvale, CA).

- 5 A preferred embodiment for amplification of a plasmid template utilizes a polymerase in the reaction mixture such that the production of amplified product does not require exogenously-added oligonucleotide primers, the amplification is exponential, and the amount of amplified product is at least 10-fold greater than the amount of plasmid template DNA put into the reaction mixture. More preferably the amplified  
10 product is at least 100-fold greater or 1,000-fold greater, even more preferably at least 10,000-fold greater or 100,000-fold greater, and most preferably at least 1,000,000-fold greater or 10,000,000-fold greater.

- The present invention also concerns the use of other reagents that further enhance amplification carried out by a core set of enzymes (the DNA polymerase, the helicase,  
15 the primase and the single-stranded binding protein). Thus, in further preferred embodiments the amplification reaction mixture also includes one or more of the following: a single-stranded binding protein of bacteriophage T7 (gene 2.5 protein), a nucleoside diphosphokinase such as that from *E. coli*, inorganic pyrophosphatase such as that from *E. coli*, an ATP regeneration system such as the combination of creatine  
20 kinase and phosphocreatine, a 5' to 3' exonuclease, preferably the 5' to 3' exonuclease is the bacteriophage T7 gene 6 exonuclease, and a ligase, preferably the ligase is the bacteriophage T7 DNA ligase. In another preferred embodiment the invention features the addition of chemical additives to the reaction mixture that increase the efficiency of the amplification reaction. Three examples of such additives are potassium glutamate,  
25 DMSO, and dextran polymer.

- In another preferred embodiment the invention features the treatment of the enzymes prior to inclusion in the reaction mixture with UV irradiation in order to reduce the amplification that arises from the contaminating DNA in the enzyme preparations. Preferably the ultraviolet light is a dose from 10 to 1000  $\mu\text{W}/\text{cm}^2$ , or preferably 100 to  
30 1000  $\mu\text{W}/\text{cm}^2$ , for from 15 sec to 15 min.

WO 00/41524

PCT/US00/00580

8

In other preferred embodiments, the amplification of the DNA is exponential, the DNA molecule to be amplified can range in size from 1,000 bp, to 2,000 bp, 3,000 bp, 5,000 bp, 10,000 bp, or even larger, up to a complete eukaryotic genome. Both single-stranded and double-stranded DNAs are amplified efficiently. When the DNA fragments  
5 being amplified are less than one thousand bases, they can be very poor substrates for amplification unless they are circular. Amplification of the DNA molecule is such that the amount of amplified product is preferably at least about 10-fold amplified, more preferably the amplified product is 100-fold greater, or at least 1000-fold greater, or at least 10,000-fold greater, or at least 100,000-fold greater, or at least 1,000,000-fold  
10 greater, or at least 10,000,000-fold greater or even more than the DNA put into the reaction mixture. The constant temperature at which the amplification reaction is carried out is preferably less than 60 °C, more preferably less than 45 °C, and more preferably at 37 °C.

In yet another preferred embodiment the reaction mixture contains ATP and CTP  
15 in addition to the 4 dNTPs.

A preferred source of the DNA molecule to be amplified is lysed *E. coli* cells and the DNA is either a plasmid, a bacterial artificial chromosome (BAC) or phage DNA. Preferably, the added plasmid DNA template is less than 100 ng and the reaction mixture is 10 to 200 µl.

20 The present method of amplification is especially useful to prepare a DNA molecule for use in a DNA sequencing reaction. Preferably, the reaction mixture containing the amplified DNA is treated with a phosphatase to remove dNTPs that may be present prior to use in a sequencing reaction. A preferred phosphatase is from arctic shrimp. The sequencing reaction preferably uses fluorescent primers or fluorescent  
25 dideoxynucleotides. A preferred embodiment for preparation of a DNA molecule for use in a sequencing reaction utilizes a polymerase in a reaction mixture such that the production of amplified product does not require exogenously-added oligonucleotide primers, the amplification is exponential, and the amount of amplified product is at least 10-fold greater than the amount of template DNA put into the mixture. More preferably  
30 the amplified product is 100-fold greater, or at least 1000-fold greater, or at least 10,000-

WO 00/41524

PCT/US00/00580

9

fold greater, or at least 100,000-fold greater, or at least 1,000,000-fold greater, or at least 10,000,000-fold greater or even more than the DNA put into the reaction mixture.

In another embodiment, the invention features a method of isolating plasmid, phage, or bacterial artificial chromosomes (BAC) from bacterial cells, such as from *E. coli* cells and then amplifying that DNA in a reaction mixture that does not require exogenous oligonucleotide primers. The bacterial cells are lysed to form a lysate which is then added to the amplification reaction mixture. Lysis can be carried out by any method known to those who practice the art, such as the use of lysozyme or detergents. A preferred embodiment for amplifying DNA directly from a bacterial cell utilizes a polymerase in a reaction mixture such that the production of amplified product does not require exogenously-added oligonucleotide primers, the amplification is exponential, and the amount of amplified product is at least 10-fold greater than the amount of DNA in the lysate that was put into the reaction mixture. More preferably the amplified product is 100-fold greater, or at least 1000-fold greater, or at least 10,000-fold greater, or at least 100,000-fold greater, or at least 1,000,000-fold greater, or at least 10,000,000-fold greater or even more.

Other types of cells can also be used. In preferred embodiments, the cells are eukaryotic microbial cells, e.g., yeast cells, fungal cells or multi-nucleate structure, or mammalian cells, e.g., human cells.

In another embodiment the invention features a method of amplifying genomic DNA in order to preserve small amounts of DNA for the purposes of long-term storage. A preferred embodiment is the amplification of human genomic DNA from a small sample from a patient (e.g. a blood sample or cheek swab) that can then be used for genotype analysis.

In another embodiment the invention features a method of determining the amount of contaminating DNA in a sample by incubating the sample potentially containing contaminating DNA with a reaction mixture comprising a DNA polymerase such that if the contaminating DNA is present the DNA molecule is amplified to produce amplified product. At least one pre-determined amount of control DNA is incubated with the same reaction mixture to produce amplified control product. The amount of amplified product in the sample is compared with the amount of amplified control DNA

WO 00/41524

PCT/US00/00580

10

as an indication of the amount of contaminating DNA in the sample. Alternatively, the rate of DNA synthesis is monitored in real-time using a probe that fluoresces only when bound to DNA; the greater the amount of input DNA, the shorter the lag period before the exponential phase of DNA synthesis can be observed (see Example 4).

- 5       Contaminating DNA refers to any DNA that may be contaminating a protein preparation. In general it is most likely to arise from the host organism in which the protein was obtained, and not purified adequately to remove it, or could be acquired during the purification of the protein from columns or reagents. Thus, the most likely source of contaminating DNA is *E. coli*, if the protein is a recombinant protein produced  
10 in *E. coli*, or it is human DNA if the protein is a recombinant protein produced in human tissue culture cells.

- By "predetermined amount of control DNA" is meant an amount of DNA such as a supercoiled plasmid (*e.g.*, pUC18 or pUC19) for example in the range of one femtogram to one nanogram in a 50 ul reaction. The synthesis obtained using this  
15 amount of DNA would be compared to an analogous reaction that was carried out in the absence of added DNA. The amount of synthesis at varying times specific for the added plasmid DNA would be determined by subtracting the amount of DNA synthesized in the absence of any DNA from that obtained in the presence of plasmid DNA. In further preferred embodiments the DNA polymerase is from bacteriophage T7, the reaction  
20 mixture further comprises a helicase, a primase and a single stranded binding protein, the control DNA is amplified at least about 10-fold, the reaction does not require exogenously-added oligonucleotide primers, the reaction mixture is carried out at a constant temperature, the control DNA is amplified at least 10-fold, and the kinetics of amplification of the control DNA is exponential.

- 25       In another aspect, the invention features a kit for amplification. The kit preferably contains a DNA polymerase, a primase, a helicase and a single-stranded binding protein. In further preferred embodiments, the components of the kit comprise those independently selected from the group consisting of a T7 DNA polymerase, the gene 4 protein from bacteriophage (the primase and the helicase), a DNA polymerase mixture of  
30 a wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced 3' to 5' exonuclease activity, and the single-stranded binding protein from *E. coli*. More

WO 00/41524

11

PCT/US00/00580

preferably, these components, a T7 DNA polymerase, the gene 4 protein from bacteriophage (the primase and the helicase), a DNA polymerase mixture of a wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced 3' to 5' exonuclease activity, and the single-stranded binding protein from *E. coli*, are selected as a group. Still more preferably, the gene 4 protein is the 63-kDa form of the protein. The kit can also contain one or more of the following components: a nucleoside diphosphokinase, an inorganic pyrophosphatase, an ATP regeneration system, preferably consisting of phosphocreatine and creatine kinase, a T7 gene 6 exonuclease, a T7 DNA ligase (gene 1.3 protein) and a single-stranded DNA binding protein of bacteriophage T7 (gene 2.5 protein).

In a further aspect the invention features a method of purifying the T7 helicase/primase (63-kDa gene 4 protein) by overproducing the protein in an *E. coli* cell preferably along with a carboxy terminal fragment of the T7 gene 4 protein, preferably the peptide comprises the carboxy terminus of the protein after residue 221 (glutamine), more preferably the peptide comprises the carboxy terminus of the protein between residues 260 (tryptophan) and residue 280 (serine), even more preferably the peptide initiates at residue 271 (arginine). This gene 4 protein peptide reduces the toxicity of the full-length gene 4 protein to the cells and increases the expression level of the full-length gene by at least 10-fold.

In another embodiment the amplification reaction is carried out in a solid matrix such as agarose. Under these conditions, in the presence of a fluorescent dye such as SYBR Green II (Molecular Probes, Inc., Eugene, OR) the amplification of single DNA molecules can be observed in a fluorescent microscope as focal centers of fluorescence that develop with time and are dependent on added DNA. In preferred embodiment, this amplification in a solid support is used to generate a library of individual "subclones" of DNA molecules, without having to passage the DNA through *E. coli* cells. More preferably these DNA clones are used as templates for DNA sequencing reactions, either by isolating individual clones and amplifying each in solution, or by carrying out the sequencing reaction using fluorescent dideoxy terminators on all the clones together simultaneously on the agarose support. This latter approach would allow a tremendous reduction in the amount (and thus the cost) of fluorescent sequencing reagents. Recently,

WO 00/41524

PCT/US00/00580

12

Mitra and Church (Nucleic Acids Research 27:e34, 1999) have described a PCR-based system to amplify and detect individual DNA molecules embedded in a polyacrylamide support.

The invention also provides a method for sequencing a DNA sequence by providing amplified DNA, that has been amplified using the constant temperature amplification method described above. The resulting products are sequenced by any of the conventional methods, such as dideoxy termination methods, preferably in an automated sequencer. Preferably the amplification is carried out in a solid matrix, preferably a gel such as an agarose or polyacrylamide gel. Preferably a circular nucleic acid construct with primer sites suitable for extension through the sequence of interest is utilized, e.g., as described in Example 10. Thus, the invention provides an efficient method of sequencing nucleic acids.

As indicated above, the source of the template sequence to be amplified can be from any of a variety of sources, including, for example, nucleic acid molecules from blood from human or non-human, cheek swabs, other tissue samples, microbial cells such as bacterial cells, viruses, plasmids or other cloning vector sequences, or other purified sequences from any source including synthetic sequences.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof and from the claims.

All articles, publications and patents cited in this application are hereby incorporated by reference, in their entirety.

#### BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 represents the nucleotide sequence of pGP4A/E-1 (SEQ. ID. NO. 1). Nucleotides 1-3 (ATG) correspond to the start codon for the 63-kDa gene 4 protein, while nucleotides 1699-1701 (TGA) correspond to its termination codon. Nucleotides 2785-2787 correspond to the start codon for the truncated gene 4 protein beginning at residue 271 (encoded for by the codon at nucleotides 2785-2787), while nucleotides 3670-3672 (TGA) correspond to the termination codon for this truncated gene.

WO 00/41524

PCT/US00/00580

13

### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following description and examples are provided for further illustrating various aspects and embodiments of the present invention and are in no way intended to be limiting in scope.

5

#### DNA polymerase

In general a polymerase for use in this invention is one that carries out extensive DNA synthesis on both strands of a DNA template, with the synthesized DNA in turn being capable of being used as a template for new DNA synthesis. This results in an exponential increase in the amount of DNA synthesized with time. Many replicative DNA polymerases have accessory proteins such as processivity factors, helicases, primases and DNA binding proteins that are specific for each DNA polymerase (for review, see Kornberg and Baker, DNA Replication, Freeman and Co., New York, 1992). Many such combinations will result in efficient DNA synthesis. The bacteriophage T7 replication system provides one example of a polymerase and accessory proteins. Those of skill in the art will appreciate that combinations of polymerases and accessory proteins from other systems (e.g. bacteriophage T4 or *E. coli*) will also be useful in the present method of amplification. Such polymerases and accessory proteins can be used with all components being from one system, or can be used in other combinations with functionally compatible proteins from two or more different systems being utilized. Those of skill in the art would be able to determine such combinations of polymerases and accessory based on the Examples and descriptions provided.

In the T7 DNA replication system, a combination of two forms of the T7 DNA polymerase result in the most efficient DNA synthesis. One is the unmodified or native T7 DNA polymerase that has high 3' to 5' exonuclease activity (Tabor *et al.*, J. Biol. Chem. 262:16212, 1987). The other is a form of T7 DNA polymerase that is deficient in its 3' to 5' exonuclease activity. An example of such a DNA polymerase is the  $\Delta 28$  T7 DNA polymerase, missing residues from lysine 118 to arginine 145 (Tabor and Richardson, J. Biol. Chem. 264:6647, 1989; U.S. Patent Nos. 4,942,130 and 4,946,786). Both these forms of T7 DNA polymerase have the processivity factor thioredoxin bound

30

WO 00/41524

PCT/US00/00580

14

to them in a one-to-one complex. DNA synthesis is most efficient when the exonuclease-deficient form is in excess over native T7 DNA polymerase; the optimum molar ratio is about 20:1. Native and exonuclease deficient polymerases are known for other replication systems, such as bacteriophage T4, bacteriophage Ø29, and *E. coli*

5 DNA polymerases I, II and III.

#### DNA helicase

Most replicative DNA polymerases require a DNA helicase for strand displacement leading strand DNA synthesis. In each replication system the helicase interacts specifically with the DNA polymerase from that system. The energy for  
10 helicase activity is obtained by the hydrolysis of nucleoside triphosphates.

The helicase of bacteriophage T7 is the gene 4 protein. Its preferred substrate for hydrolysis is dTTP. The phage makes two forms of the gene 4 protein of molecular weight 56,000 and 63,000; the two forms arise from two in-frame start codons. Efficient amplification requires the 63-kDa form of the gene 4 protein, since, as discussed below,  
15 this form also provides primase activity (Bernstein and Richardson, *J. Biol. Chem.* 264:13066, 1989). Altered forms, substitutions, insertions, deletions, of the 63-kDa protein are also suitable for the present invention. One example is the 63-kDa gene 4 protein in which the methionine at residue 64 is changed to a glycine (G4A<sub>G64</sub>). (Mendelman *et al.*, *Proc. Natl. Acad. Sci. USA* 89:10638, 1992; Mendelman *et al.*,  
20 *J. Biol. Chem.* 268:27208, 1993). This form of the 63-kDa gene 4 protein is useful because the mutation prevents the initiation of synthesis of the 56-kDa gene 4 protein. All enzymatic properties of the G4A<sub>G64</sub> form of the gene 4 protein that have been examined are comparable to those of the wild-type 63-kDa gene 4 protein, including its use as a primase and helicase for amplification as described in the current invention.

25

#### DNA primase

In most replication systems, synthesis of the lagging strand is initiated from short oligoribonucleotide primers that are synthesized at specific sites by primases. Specific interactions between a primase and the DNA polymerase allow the DNA polymerase to initiate DNA synthesis from the oligoribonucleotide resulting in the synthesis of the  
30 lagging strand.



WO 00/41524

15

PCT/US00/00580

A unique primase mechanism is found in the terminal protein exemplified by adenovirus and bacteriophage  $\phi$ 29 (Salas, Ann. Rev. Biochem. 60:39, 1991). In these organisms, duplex linear DNA is replicated by the covalent binding of the terminal protein to each 5' end, followed by initiation of DNA synthesis from a mononucleotide bound to the terminal protein by a DNA polymerase. While this system can provide a very efficient form of amplification of the phage genomes such as  $\phi$ 29 DNA (Blanco *et al.*, Proc. Natl. Acad. Sci. USA 91:12198, 1994), the nature of this priming mechanism makes it a difficult system to modify for general amplification of DNAs.

In the T7 replication system, the gene 4 protein, in addition to being a helicase, is also the primase (Bernstein and Richardson, Proc. Natl. Acad. USA 85:396, 1988; Bernstein and Richardson, J. Biol. Chem. 264:13066, 1989). Only the 63-kDa form of the gene 4 protein has primase activity. At specific pentanucleotide recognition sites, the 63-kDa gene 4 protein synthesizes tetranucleotides that are used as primers by T7 DNA polymerase to initiate lagging strand DNA synthesis. The synthesis of RNA primers by 63-kDa gene 4 protein requires ATP and CTP. However, the addition of exogenous short oligonucleotides (7 bases or less) can be used by the gene 4 protein to initiate DNA synthesis primers by T7 DNA polymerase (Scherzinger *et al.*, Nucleic Acids Research 4:4151, 1977). The 63-kDa gene 4 protein is essential for amplification using the T7 replication system. While amplification is most efficient when ATP and CTP are present, it is possible that short oligonucleotides could be added to the reaction mixture to provide priming for amplification in the absence of ATP and CTP. While the native T7 DNA polymerase is capable of interacting with the 63-kDa gene 4 protein to initiate DNA synthesis from RNA primers, the  $\Delta$ 28 T7 DNA polymerase (missing residues 118 to 145) does so much less efficiently. This may be one explanation why a small amount of native T7 DNA polymerase is critical for efficient amplification.

#### Single-stranded DNA binding protein

Most replication systems require single-stranded DNA binding proteins. They serve a number of roles (for review, see Kornberg and Baker, DNA Replication, Freeman and Co., New York, 1992). For example, they remove secondary structure from single-stranded DNA to allow efficient DNA synthesis.

WO 00/41524

PCT/US00/00580

16

The single-stranded DNA binding protein (ssb) from *E. coli* has been extensively characterized. It has a dramatic effect on improving the processivity of T7 DNA polymerase, particularly at temperatures below 30° C (Tabor *et al.*, J. Biol. Chem. 262:16212, 1987). A low level of ssb is required for efficient amplification using the T7 replication proteins described in this invention. Optimally, the amount of ssb for a 50 µl reaction is from 0.01 to 1 µg.

Bacteriophage T7 also encodes its own single-stranded DNA binding protein, the gene 2.5 protein (Kim *et al.*, J. Biol. Chem. 267:15022, 1992). The presence of this protein is less important for amplification than is the *E. coli* single-stranded DNA binding protein. However, it does stimulate the rate of DNA synthesis several-fold at a concentration of from 0.05 to 5 µg in a 50 µl reaction containing the core amplification components.

#### Nucleoside Diphosphokinase

Nucleoside diphosphokinase rapidly transfers the terminal phosphate from a nucleoside triphosphate to a nucleoside diphosphate. It is relatively nonspecific for the nucleoside, recognizing all four ribo- and deoxyribonucleosides. Thus it efficiently equilibrates the ratio of nucleoside diphosphates and nucleoside triphosphates among all the nucleotides in the mixture. This enzyme can increase the amount of DNA synthesis if one of the required nucleoside triphosphates is preferentially hydrolyzed during the reaction.

In the bacteriophage T7 replication system, the T7 helicase (gene 4 protein) hydrolyzes dTTP to dTDP to provide energy (Matson and Richardson, J. Biol. Chem. 258:14009, 1983). Thus the concentration of dTTP will decrease disproportionately during DNA synthesis. The addition of nucleoside diphosphokinase will re-equilibrate the level of triphosphates of all the nucleosides present. This results in an increase of several-fold in the amount of DNA synthesized. Nucleoside diphosphokinase from Baker's Yeast can be purchased from Sigma Chemical Co. (St. Louis, MO) and the overproduction and purification of nucleoside diphosphokinase from *E. coli* has been described (Almaula *et al.* J. Bact. 177:2524, 1995). Other nucleoside diphosphokinases are known to those who practice the art and are useful in the present invention.

WO 00/41524

PCT/US00/00580

17

#### Inorganic Pyrophosphatase

During a DNA synthesis reaction, inorganic pyrophosphate will accumulate as a product of the reaction. If the concentration becomes too high, it can reduce the amount of DNA synthesis due to product inhibition. This can be prevented by the addition of  
5 inorganic pyrophosphatase. Yeast inorganic pyrophosphatase can be purchased from Sigma Chemical Co. (St. Louis, MO). Other inorganic pyrophosphatases are known to those who practice the art and are useful in the present invention.

#### ATP-regenerating system

During DNA synthesis in many DNA replication systems, some of the  
10 deoxynucleoside triphosphates will be degraded to deoxynucleoside diphosphates due to hydrolysis by the helicase present. This can be minimized by the use of an ATP regeneration system which, in the presence of nucleoside diphosphokinase, will convert any nucleoside diphosphate in the reaction mixture to the triphosphate. In the T7 replication system, the helicase very rapidly degrades dTTP to dTDP for energy. The  
15 presence of an ATP-regeneration system will increase the amount of nucleotides capable of serving as precursors for DNA synthesis.

A number of ATP regeneration systems are available and have been widely used. For example, the combination of phosphocreatine and creatine kinase will push the equilibrium between ADP and ATP towards ATP, at the expense of the phosphocreatine.  
20 Both phosphocreatine and creatine kinase are available from Sigma Chemical Co. (St. Louis, MO).

#### Joining lagging strand fragments

In most DNA replication systems, the lagging strand is synthesized as a series of short, "Okazaki" fragments, that are initiated by a short RNA primer at each 5' end. In  
25 order to form a continuous strand, a 5' to 3' exonuclease is needed to remove the RNA primer, and then, after a DNA polymerase fills in the gap, a DNA ligase is needed to seal the nick. In the T7 DNA replication system, these functions are accomplished by the action of the gene 6 protein, a 5' to 3' exonuclease, and the gene 1.3 protein, a DNA ligase (Engler and Richardson, J. Biol. Chem. 258:11197, 1983). Purification of the  
30 gene 6 protein (Kerr and Sadowski, J. Biol. Chem. 247:305, 1972; Engler and

WO 00/41524

PCT/US00/00580

18

Richardson, *J. Biol. Chem.* 258:11197, 1983) and gene 1.3 protein (Doherty *et al.*, *J. Biol. Chem.* 271:11083, 1996) have been described. Exonucleases and ligases from other replication systems are known to those who practice the art and are useful in the present invention.

5

#### Removal of contaminating DNA from enzymes

Since the amplification system described in this invention is nonspecific, contaminating DNA will potentially be a serious source of background. This is particularly a problem in enzyme preparations in which the enzymes bind DNA, such as DNA polymerases, helicases and DNA binding proteins. Most DNA can be removed from enzyme preparations during purification. For example, an anion exchange column such as DEAE will bind DNA much tighter than it binds proteins.

The residual DNA present in enzyme preparations can be inactivated as a template for DNA synthesis by treatment of the enzymes with ultraviolet light. Effective doses of ultraviolet light are 10 to 1000  $\mu\text{W}/\text{cm}^2$ . In the T7 replication system, treatment of the enzymes with an ultraviolet dose of 200  $\mu\text{W}/\text{cm}^2$  for 2 min reduces the background synthesis due to contaminating DNA by 99%, while it reduces the activity of the enzymes by less than 20%. This dose of ultraviolet light is equivalent to the dose obtained using a standard, hand-held short-wave ultraviolet light source 10 cm from the enzyme mixture. The enzyme mixture is kept on ice during the treatment with ultraviolet light to help maintain its activity.

20

#### Removal of unincorporated nucleoside triphosphates from reaction mixture

In order to be used for DNA sequencing reactions, the template DNA added to the reactions can only have low levels of unincorporated dNTPs. An efficient method of removing dNTPs is to treat the amplified product with alkaline phosphatase such as that from arctic shrimp (Amersham Pharmacia Biotech, Inc., Piscataway, NJ). This phosphatase is rapidly inactivated by treatment at elevated temperatures. Thus treatment of a 50  $\mu\text{l}$  reaction mixture with 0.1 to 10 units of shrimp alkaline phosphatase at 20-37°C for 10 to 60 min, followed by treatment at 65-95°C for 10 to 60 min, results in template DNA free of dNTPs that can be added directly to a DNA sequencing reaction such as the dye terminator reactions using AmpliTaq FS enzyme (Applied Biosystems,

30

WO 00/41524

PCT/US00/00580

19

Foster City, CA) or ThermoSequenase enzyme (Amersham Pharmacia Biotech Inc., Piscataway, N.J).

Reaction mixture for amplification using T7 replication proteins

Reaction mixtures contain a buffer at a concentration of 5 mM to 50 mM at pH 7.0 to 8.5, preferably pH 7.5; e.g., 20-40 mM Tris-glutamate, pH 7.5. 2 mM to 15 mM  $Mg^{2+}$  is required, preferably 5 to 10 mM; e.g., 9 mM  $MgCl_2$ . A reducing agent is present, such as 1 to 50 mM dithiothreitol or 2-mercaptoethanol; e.g. 5 mM dithiothreitol. The reaction mixture contains 50 to 200 mM potassium glutamate; e.g. 100 mM potassium glutamate, which stimulates the reaction. In contrast chloride salts generally inhibit the reaction; if NaCl or KCl are present, preferably their concentration is below 50 mM. The 4 dNTPs are present at a concentration preferably between 20 and 2,000  $\mu M$ , most preferably at a concentration of 500-1000  $\mu M$  of each dNTP. rATP and rCTP are each present at a concentration between 20 and 2000  $\mu M$ , most preferably at a concentration of 500 to 1000  $\mu M$ . Other chemicals may be added that enhance the amplification reaction. For example, DMSO between 0.5 and 8% gives an increased amount of DNA synthesis, most preferably a concentration of 4%. Similarly, a dextran such as Dextran T-10 or Dextran T-500 polysaccharide (Amersham Pharmacia Biotech Inc., Piscataway, NJ) between 0.5 and 20% enhances the amount of DNA synthesis, most preferably a concentration of 5%. After the addition of the DNA to be amplified and the enzyme mixture containing at least some of the T7 replication enzymes described above, the reaction is allowed to proceed at a temperature between 10 ° C and 50° C, preferably 37° C, for 10 to 60 min, preferably 20 min. The reaction can be stopped using standard procedures such as the addition of EDTA to a final concentration of 25 mM, or heating the mixture at 70° C for 20 minutes.

25      Sequencing from Single Molecule Amplification

The amplification reaction can be used to provide DNA for sequencing. The amplification is carried out in a solid matrix such as agarose. Under these conditions, in the presence of a fluorescent dye such as SYBR Green II (Molecular Probes, Inc., Eugene, OR) the amplification of single DNA molecules can be observed in a fluorescent microscope as focal centers of fluorescence that develop with time and are

30

WO 00/41524

PCT/US00/00580

20

dependent on added DNA. Low-melt agarose such as SeaPlaque Agarose (FMC Products, Rockland, ME) is preferred as the matrix, since the DNA can be easily removed from the agarose when desired by heat. It may be helpful to use a combination of agaroses or other matrices to optimize the stability of the individual foci in the solid support while allowing the DNA to be readily extracted when desired.

If the single molecules being amplified are from a large DNA fragment that has been digested with a restriction enzyme (e.g. EcoRI), and then ligated intramolecularly to form circular molecules, the collection of foci within the agarose support will correspond to a population of "subclones" of that original DNA molecule, comparable to subclones obtained more conventionally by transforming these molecules into *E. coli* and then plating the cells to obtain individual clones. The advantage of the approach described here is that the entire library of clones is generated without the need to transform the DNA into bacteria, and that each "clone" observed as a foci in agarose is pure DNA that can be used directly for sequencing or other applications. Recently, Mitra and Church (Nucleic Acids Research 27:e34, 1999) have described a PCR-based system to amplify and detect individual DNA molecules embedded in a polyacrylamide support.

Several different approaches can be used to sequence the DNA amplified in the agarose support. One would be to pick the DNA from each foci into a tube, heat at 70 °C to melt the agarose, and then add an aliquot to a new amplification reaction mixture to amplify the DNA in solution. This amplified DNA could then be used for DNA sequence analysis as described in Example 5.

Alternatively, the DNA on the agarose support could be sequenced directly. The unincorporated nucleotides from the amplification would first be removed by soaking the agarose in a suitable buffer, e.g., 10 mM Tris-HCl, pH 7.5, 1 mM EDTA. The DNA in the agarose would then be denatured by treatment with 10 mM NaOH for 5 min, followed by neutralization by several washes in 10 mM Tris-HCl, pH 7.5, 1 mM EDTA. The primer used for DNA sequencing would then be soaked into the agarose and allowed to anneal to the denatured DNA. Excess primer would be removed by several washes in 10 mM Tris-HCl, pH 7.5, 1 mM EDTA, and then the DNA sequencing reactions would be carried out by the addition of sequencing reaction buffer, fluorescent nucleotides (e.g. BigDye nucleotides from PE Biosystems (Foster City, CA)), and a DNA polymerase

WO 00/41524

PCT/US00/00580

21

such as AmpliTaq FS (PE Biosystems, Foster City, CA). The reaction would be initiated by heating the sample to 70 °C. After a 10 min reaction, the sample would be chilled to 4 °C, and the product DNA would be denatured by treatment again with 10 mM NaOH for 5 min. Finally, the salt would be removed by extensive washing in water.

5       The samples are now ready to be directly loaded onto a capillary DNA sequencing instrument. Injection of the capillaries would be a modification of that used in existing instruments such as the PEBiosystems 3700 Genetic Analyzer (Foster City, CA). One direct-load approach would be to insert a 100-1000 capillary array into the sample, heat to 70 °C to melt the agarose, and then apply a voltage to electrosmotically  
10       inject the DNA into each capillary. While only a subset of the capillaries would contain DNA from a single foci, and thus produce unique DNA sequence, the advantages of this procedure would be the ease of carrying out multiple sequencing reactions simultaneously and the low reagent cost of carrying out such parallel reactions.

15

Example 1: Amplification of purified plasmid DNA using a minimal number of T7 replication proteins

The reaction mixture (45 µl) contains 20 mM Tris-glutamate, pH 7.5, 9 mM MgCl<sub>2</sub>, 6 mM dithiothreitol, 100 mM potassium glutamate, 3.5% dimethylsulfoxide  
20       (DMSO), 7% Dextran T-500 polysaccharide (Amersham Pharmacia Biotech, Inc., Piscataway, NJ), 550 µM each dGTP, dATP, dTTP and dCTP, 330 µM ATP, 440 µM CTP, and DNA. The DNA samples range from 0.1 pg to 10 ng of supercoiled double-stranded plasmid DNA.

Although circular DNA is a preferred substrate for amplification, linear DNA can  
25       also be used. The DNA can be either single- or double-stranded. In addition to plasmid DNA, other DNAs such as phage lambda DNA, phage M13 DNA bacterial artificial chromosomal (BAC) DNA and genomic DNA (*e.g.*, bacterial or human) are effective templates for amplification. Synthetic DNA such as oligonucleotides 100 nucleotides in length can be used if they are first ligated to form circular molecules. Data suggest that

WO 00/41524

PCT/US00/00580

22

for circular DNA molecules the initial stages of amplification involve a rolling circle mechanism.

One method of initiating rolling circle DNA synthesis on a plasmid DNA from a specific site that has been described is to use the site-specific single-stranded  
5 endonuclease gene II protein from filamentous phage such as M13 or fd. This protein recognizes a sequence of approximately 50 bp (the plus strand replication origin) and introduces a specific nick into one of the strands (Higashitani *et al.*, Nucleic Acids Research 20:2685, 1992). While the wild-type gene II protein requires supercoiled DNA for activity, mutant proteins have been described (*e.g.* glycine 73 changed to alanine) that  
10 reduce this superhelicity requirement, so that the protein now recognizes relaxed duplex DNA (Higashitani *et al.*, Nucleic Acids Research 20:2685, 1992). The purification of the wild-type and mutant gene II proteins have been described (Greenstein and Horiuchi, J. Molec. Biol. 197:157, 1987; Higashitani *et al.*, Nucleic Acids Research 20:2685, 1992). It has been demonstrated that cleavage of supercoiled DNA with the gene II  
15 protein results in a substrate that the T7 DNA polymerase and T7 gene 4 protein can use to initiate lagging strand DNA synthesis (Harth *et al.*, Eur. J. Biochem. 119:663, 1981). It has also been demonstrated that these molecules promote rolling circle DNA synthesis using the replication proteins from bacteriophage T4 (Meyer *et al.*, J. Biol. Chem. 256:5810, 1981). This class of site-specific nicking proteins may be useful for  
20 stimulating initiation of the amplification process described in this invention. However, in the invention described here the inventors have shown that surprisingly nicking of the circular DNA is not necessary for efficient initiation of DNA synthesis; the combination of T7 gene 4 protein and T7 DNA polymerase are able to efficiently initiate and synthesize DNA on covalently supercoiled plasmid DNAs whether they are supercoiled  
25 or relaxed.

For diagnostic characterization of the extent of DNA synthesis, one of the dNTPs can be radioactively labeled; for example, [<sup>3</sup>H] dTTP can be used at a specific activity of 20 cpm/pmol. The synthesized DNA can also be detected by fluorescence or chemiluminescence by incorporating the appropriate modified nucleotides using standard  
30 techniques (Current Protocols in Molecular Biology, Ausubel *et al.*, eds., John Wiley &



WO 00/41524

PCT/US00/00580

23

Sons, Inc., 1996). ). In addition, the rate of DNA synthesis can be monitored in real time by the addition of a fluorescent probe to the reaction mixture (see Example 4).

The enzyme mixture (5  $\mu$ l per reaction) is prepared in a buffer containing 20 mM Tris-glutamate, pH 7.5, 10 mM 2-mercaptoethanol and 0.5 mg/ml bovine serum albumin.

- 5 5  $\mu$ l of this mixture contains 1  $\mu$ g of  $\Delta$ 28 T7 DNA polymerase, 0.03  $\mu$ g of native T7 DNA polymerase, 1  $\mu$ g of T7 63-kDa gene 4 protein, and 0.3  $\mu$ g of *E. coli* single-stranded DNA binding protein.  $\Delta$ 28 T7 DNA polymerase is a one-to-one mixture of *E. coli* thioredoxin and T7 gene 5 protein with a deletion of residues 118 to 145, as described in Tabor and Richardson, J. Biol. Chem. 264:6647, 1989, and U.S. Patents
- 10 Nos. 4,942,130 and 4,946,786. It is the same as Sequenase Version 2.0 enzyme sold by Amersham Pharmacia Biotech, Inc. (Piscataway, NJ). Native T7 DNA polymerase is a one-to-one complex of thioredoxin and wild-type T7 gene 5 protein. It can be purchased from Amersham Pharmacia Biotech, Inc. (Piscataway, NJ) or New England Biolabs (Beverly, MA). T7 63-kDa gene 4 protein is the G4A<sub>G64</sub> protein described in
- 15 Mendelman *et al.*, Proc. Natl. Acad. Sci. USA 89:10638, 1992 and Mendelman *et al.*, J. Biol. Chem. 268:27208, 1993; it is the wild-type gene 4 63 kDa protein except that the methionine at residue 64 has been replaced with a glycine to prevent initiation of synthesis of the 56-kDa form of the gene 4 protein. *E. coli* single-stranded DNA binding protein (ssb) is overproduced and purified as described in Lohman *et al.*, Biochemistry
- 20 25:21 (1986). It can be purchased from Amersham Pharmacia Biotech, Inc. (Piscataway, NJ).

- In order to reduce the level of background DNA synthesis resulting from contaminating DNA, the enzyme mixture is treated with ultraviolet light at a dose of 200  $\mu$ W/cm<sup>2</sup> for 2 min on ice prior to adding to the reaction mixture. This dose corresponds
- 25 to a distance of about 10 cm from a standard short-wave ultraviolet lamp. Times of 10 sec to 15 min are also effective at reducing the amount of contaminating DNA that can be replicated without affecting significantly the activity of the enzymes.

To initiate the amplification reaction, 5  $\mu$ l of the enzyme mixture is added to the 45  $\mu$ l reaction mixture. The reaction is then allowed to proceed at 37° C for 20 min. The

WO 00/41524

PCT/US00/00580

24

reaction can be carried out at a temperature of 10°C to 45°C, for 1 min to 2 hr. The reaction is stopped by the addition of 5 µl of 200 mM EDTA.

There are many published procedures to measure the amount of DNA synthesized during the reaction. For example, if [<sup>3</sup>H]dTTP is present, the amount of radioactivity incorporated into DNA can be measured by binding the reaction mixture to a DE81 filter disk, then washing the disk with 0.3 M ammonium formate to remove unincorporated nucleotides, and then measuring the amount of radioactivity retained on the disk using a scintillation counter, as described in Kim *et al.*, *J. Biol. Chem.* 267:15032 (1992). Alternatively, the DNA can be analyzed by agarose gel electrophoresis and staining with ethidium bromide, either with or without digestion with restriction enzymes, following standard procedures (Current Protocols in Molecular Biology, Ausubel *et al.*, eds., John Wiley & Sons, Inc., 1996).

This procedure should result in the incorporation of about 5% of the total dNTPs into DNA, or about 1.5 µg of DNA synthesized in the 50 µl reaction mixture. If the reaction mixture contains 10 pg of plasmid DNA, this corresponds to a 150,000-fold amplification.

**Example 2: Amplification of purified plasmid DNA using a T7 replication system optimized for maximum rate and amount of DNA synthesis**

The rate and amount of DNA synthesis obtained using the conditions defined in Example 1 can be improved by using the modifications described below. All other components and conditions are identical to those outlined in Example 1.

Phosphocreatine (11 mM) is added to the 45 µl reaction mixture. This in combination with creatine kinase provides an efficient ATP regeneration system. The phosphocreatine that is most effective is synthetic phosphocreatine, Sigma catalog number P6502 (Sigma Chemical Co., St. Louis, MO), dissolved in H<sub>2</sub>O to a concentration of 500 mM.

The enzyme mixture contains at least some of the following six enzymes. These are in addition to the Δ28 T7 DNA polymerase, native T7 DNA polymerase, 63-kDa

WO 00/41524

PCT/US00/00580

25

gene 4 protein and *E. coli* single-stranded DNA binding protein, which are present in the amounts defined in Example 1.

(1) Creatine kinase, rabbit muscle (Boehringer catalog number 127566) (Boehringer Mannheim, Indianapolis, IN). A stock solution is prepared at 100 mg/ml in H<sub>2</sub>O and stored at -40° C. Each enzyme mixture (5 µl) for a 1X reaction (50 µl total volume) contains 2 µg of creatine kinase. Creatine kinase, in conjunction with phosphocreatine and nucleoside diphosphokinase, provides an ATP regeneration system that converts nucleoside diphosphates that arise during the reaction to nucleoside triphosphates.

(2) Nucleoside diphosphokinase, Baker's yeast (Sigma catalog number N0379) (Sigma Chemical Co., St. Louis, MO). A stock solution is prepared at 1 mg/ml in H<sub>2</sub>O and stored at -40° C. Each enzyme mixture (5 µl) for a 1X reaction (50 µl total volume) contains 50 ng of nucleoside diphosphokinase. Alternatively, nucleoside diphosphokinase can be overproduced and purified from *E. coli* (Almaula *et al.*, *J. Bact.* 177:2524, 1995). 50 ng of the *E. coli* enzyme is used for a 1X reaction. Nucleoside diphosphokinase is added to maintain an equal ratio of all four dNTPs.

(3) Inorganic pyrophosphatase, Baker's yeast (Sigma catalog number I1891) (Sigma Chemical Co., St. Louis, MO). A stock solution is prepared at 1 mg/ml in H<sub>2</sub>O and stored at -40° C. Each enzyme mixture (5 µl) for a 1X reaction (50 µl total volume) contains 20 ng of inorganic pyrophosphatase. Inorganic pyrophosphatase is added to degrade the pyrophosphate that will build up during polymerization and could inhibit DNA synthesis by product inhibition.

(4) T7 single-stranded DNA binding protein (T7 gene 2.5 protein). T7 single-stranded DNA binding protein is overproduced and purified as described by Kim *et al.* (*J. Biol. Chem.*, 267:15022, 1992). Each enzyme mixture (5 µl) for a 1X reaction (50 µl total volume) contains 1 µg of T7 single-stranded DNA binding protein. The T7 single-stranded DNA binding protein will increase the rate of DNA synthesis by several-fold. One possible explanation for this is that the T7 single-stranded DNA binding protein is necessary to have efficient coupling of leading and lagging strand DNA synthesis by the

WO 00/41524

PCT/US00/00580

26

T7 DNA polymerase and helicase/primase complex (Park *et al.*, J. Biol. Chem. 273:5260, 1998; Lee *et al.*, Mol. Cell. 1:1001, 1998).

(5) T7 gene 6 exonuclease. The combination of T7 gene 6 exonuclease and T7 DNA ligase (T7 gene 1.3 protein) are used to remove the RNA primers from the 5' ends of the lagging strand fragments and then, after the gaps are filled in by the wild-type T7 DNA polymerase, seal the resulting nicks. The T7 gene 6 exonuclease is purified as described in Kerr and Sadowski, J. Biol. Chem. 247:305, 1972 and Engler and Richardson, J. Biol. Chem. 258:11197, 1983. Each enzyme mixture (5  $\mu$ l) for a 1X reaction (50  $\mu$ l total volume) contains 50 ng of T7 gene 6 exonuclease.

(6) T7 DNA ligase (gene 1.3 protein). The T7 ligase is added to seal any nicks that are present in the synthesized DNA, in particular those that occur on the lagging strand. T7 DNA ligase is overproduced and purified as described in Doherty *et al.*, J. Biol. Chem. 271:11083, 1996. Each enzyme mixture (5  $\mu$ l) for a 1X reaction (50  $\mu$ l total volume) contains 0.1  $\mu$ g of T7 DNA ligase.

As in Example 1, the enzyme mixture is treated with an ultraviolet dose of 200  $\mu$ W/cm<sup>2</sup> for 2 min on ice prior to adding to the reaction mixture. The amplification reaction is then initiated by the addition of 5  $\mu$ l of the enzyme mixture to the 45  $\mu$ l reaction mixture. The reaction is allowed to proceed at 37° C for 20 min, and then stopped by the addition of 5  $\mu$ l of 200 mM EDTA. The products are analyzed as described in Example 1.

This procedure should result in the incorporation of up to 50% of the total dNTPs into DNA, or about 15  $\mu$ g of DNA synthesized in the 50  $\mu$ l reaction mixture. If the initial reaction mixture contains 10 pg of plasmid DNA, this corresponds to a 1,500,000-fold amplification.

Example 3: Assay to determine whether amplification is exponential

Reactions as described in Examples 1 and 2 are carried out using varying amounts of plasmid DNA and for varying lengths of time. A radioactively labeled dNTP is used in order to determine the amount of DNA synthesis; for example, 50 cpm/pmole [<sup>3</sup>H]dTTP or [ $\alpha$ -<sup>32</sup>P]dATP. 100  $\mu$ l reactions are carried out in the absence of added

WO 00/41524

27

PCT/US00/00580

DNA, and in the presence of 1, 10, 100 and 1,000 pg of supercoiled pUC18 plasmid DNA. Reactions are carried out at 37° C and 20 µl aliquots are removed at 5, 10, 15, 20 and 30 min and stopped by the addition of 5 µl of 200 mM EDTA, pH 8.0. The amount of radioactivity incorporated into DNA is determined using standard methods. For example, the amount of radioactivity that can be precipitated by trichloroacetic acid, which will precipitate DNA but not free nucleotide, can be determined as described in Tabor and Richardson, *J. Biol. Chem.* 264:6647, 1989. Alternatively, the amount of radioactivity that is retained by DE81 filter disks in the presence of 0.3 M ammonium formate, pH 8.0, can be determined (Kim *et al.*, *J. Biol. Chem.* 267:15032, 1992); DNA but not dNTPs is retained by the disks under these conditions.

For each time point, the amount of DNA synthesized in the absence of added DNA is subtracted from the amount synthesized in the presence of DNA. Up to 20 min, the amount of DNA synthesized in the absence of added input DNA should be very low (for example, less than 1% the amount synthesized in the presence of 100 pg of added input DNA).

For each amount of input DNA, the amount of DNA synthesized above the background level synthesized in the absence of input DNA is plotted as a function of the time of the reaction. If amplification is exponential, then the rate of synthesis of DNA will increase during some portion of the time course (i.e., the plot will be sigmoidal). For example, the amount of DNA synthesized after 15 min might be 50-fold higher than the amount synthesized after 10 min.

#### Example 4: Real-time amplification assay using fluorescence

A very effective method of monitoring the amount of DNA synthesized using the isothermal exponential amplification system is to use fluorescent probes to continuously report the amount of DNA in the reaction in real time. The basic principle of this method has been used successfully to determine the initial concentrations of specific RNAs and DNAs in a PCR reaction (Gibson *et al.*, *Genome Research* 6:995, 1996). The higher the initial DNA concentration, the shorter the lag period before the exponential phase of DNA synthesis. To measure the amount of DNA synthesis in real time, reactions were carried out as described in Example 3 except that radioactivity was

WO 00/41524

28

PCT/US00/00580

omitted and the fluorescent dye SYBR Green II (Molecular Probes, Eugene, OR) was added at a final concentration of 1/80,000 the stock solution. The dye was added after the reaction mixture was treated with UV light to destroy contaminating DNA. SYBR Green II has virtually no fluorescence on its own but fluoresces very strongly when  
5 bound to nucleic acid. At concentrations higher than 1/80,000 of the initial stock it is inhibitory to the amplification reaction described in Example 3. Other dyes that fluoresce specifically when bound to nucleic acid will also work in the present invention (e.g. SYBR Green I, Pico Green, Oligreen, SYTO 11, SYTO 12, SYTO 13, SYTO 14, SYTO 15, SYTO 16 (Molecular Probes, Eugene, OR), and SYBR Green (PE  
10 Biosystems, Foster City, CA)) or any other dye which shows significantly enhanced fluorescence when bound to DNA. Preferably, the significantly enhanced fluorescence refers to at least a 10-fold increase in fluorescence upon binding to DNA. More preferably, DNA binding will result in at least a 30-fold, 100-fold, 300-fold or 1000-fold increase in fluorescence.

15 In the example, reactions are carried out at 37 °C. Fluorescence is monitored once every 15 sec for 30 min. Exemplary instruments that can be used to measure fluorescence include real-time PCR instruments such as the PE Biosystems 5700 (PE Biosystems, Foster City, CA) and the Roche LightCycler (Indianapolis, IN), and fluorescence microtiter plate readers that can maintain a constant temperature and can  
20 carry out kinetic measurements, such as the SPECTRAmax (Molecular Devices, Sunnyvale, CA). Ideally the SYBR Green II fluorescence is monitored using an excitation wavelength of 480 nM and an emission wavelength of 530 nM.

In order to measure the DNA concentration in unknown samples, a standard curve is determined for each experiment by carrying out reactions either in the absence  
25 of DNA (blank) or in the presence of known concentrations of a standard DNA such as supercoiled pUC18 (e. g. ten-fold dilutions from 0.1 pg to 10 ng). A standard curve is determined by plotting the time required for fluorescence to reach half-maximum (or alternatively the time required for the relative fluorescence to reach some threshold value) versus the log of the initial concentration of DNA. This plot yields a straight line  
30 over at least three orders of magnitude of initial DNA concentration. A comparison of

WO 00/41524

PCT/US00/00580

29

the values obtained using unknown samples to the standard curve allows one to extrapolate the amount of DNA present at the outset in each sample.

Example 5: Use of amplified DNA as a template for DNA sequencing reactions

5 The following example shows how the amplification reaction can be used to generate enough template from a small amount of DNA to be sequenced using conventional radioactive and nonradioactive dideoxy terminator sequencing strategies. While the example shown is starting from a small amount of plasmid DNA, this procedure will also work using other DNAs such as those from bacterial artificial  
10 chromosomes (BACs), phage lambda, phage M13, or genomic DNA (*e.g.*, bacterial or human). Also, while the example shown uses purified DNA, much less pure samples, such as that from freshly lysed *E. coli* cells, can also be used.

Supercoiled pUC18 DNA (100 pg) is added to a 50  $\mu$ l reaction mixture as described in Example 2, and amplified at 37°C for 20 min. The reaction is stopped by  
15 heating the mixture at 70°C for 5 min. 0.5 units of arctic shrimp alkaline phosphatase (Amersham catalog number E70092) (Amersham Pharmacia Biotech, Inc., Piscataway, NJ) is added, and the mixture is incubated at 37°C for 30 min. The phosphatase is then inactivated by incubation at 80°C for 15 min. The amplified DNA can now be directly added to a DNA sequencing reaction. Other methods can also be used to purify the  
20 template DNA away from unincorporated dNTPs. These include precipitating the DNA with ethanol or isopropanol, and purifying the DNA by binding it to a silica matrix, for example using a commercial kit such as the QIAquick PCR Purification Kit, product number 28104 (Qiagen, Inc., Valencia, CA).

In order to sequence the amplified DNA using the PE Biosystems BigDye  
25 Terminators (catalog number 4303149 (PE Biosystems, Foster City, CA), 1  $\mu$ l of the amplified DNA (500 ng) that had been treated with arctic shrimp phosphatase was added to 8  $\mu$ l of the BigDye Reaction Mixture, 1.5  $\mu$ l (3 pmoles) of the "universal" forward primer, and 9.5  $\mu$ l of water. Cycle sequencing was carried out using 25 cycles consisting of 96 °C for 10 sec, 50 °C for 5 sec, and then 60 °C for 4 min. After the reaction was  
30 complete the unincorporated dye terminators were removed by gel filtration centrifugation using Centri-Sep columns (Princeton Separations, Inc., Princeton, NJ) and

WO 00/41524

PCT/US00/00580

30

the eluent was used directly for capillary sequence analysis on the PE Biosystems 310 single-capillary Genetic Analyzer or the PE Biosystems 3700 96-capillary Genetic Analyzer. The accuracy of the sequence was analyzed by determining the number of bases that had a Phred score greater than 20 (Ewing et al., Genome Research 8:186, 5 1998). In general, three times less amplified DNA was required compared with supercoiled plasmid DNA to produce DNA sequence of a given quality as analyzed by their relative Phred scores. This is probably due to the fact that primers will anneal more efficiently to the amplified DNA since it is linear compared with the supercoiled plasmid DNA.

10 The amplified DNA can also be sequenced using other methods of labeling the fragments. For example, it can be used a template for reactions containing radioactively labeled dideoxyterminators; 2  $\mu$ l of the DNA can be added directly to the 20  $\mu$ l reaction mixture from a ThermoSequenase radiolabeled terminator cycle sequencing kit, Amersham Pharmacia product number US 79750 (Amersham Pharmacia Biotech, Inc., 15 Piscataway, NJ). The DNA can also be used for DNA sequencing techniques that use either radioactive or fluorescently labeled primers.

Example 6: Amplification and sequencing of plasmid DNA directly from a bacterial colony

20 The procedure below describes the amplification *in vitro* of DNA obtained from direct lysis of a bacterial colony, followed by DNA sequencing of the amplified DNA. While the example is using cells containing a high copy plasmid (pUC18), one skilled in the art will recognize that cells containing other types of DNAs, such as bacterial artificial chromosome (BAC) DNAs, could also be used in this procedure.

25 Bacterial cells containing the plasmid to be amplified and sequenced, for example a pUC18 derivative, are plated on LB plates containing 50  $\mu$ g/ml ampicillin at 37° C overnight. A single colony is placed in 500  $\mu$ l of a solution containing 20 mM Tris-HCl, pH 7.5, 1 mM EDTA and 100  $\mu$ g lysozyme. After incubation at room temperature for 15 min, the solution is centrifuged at 14,000 rpm for 10 min. 1  $\mu$ l of the supernatant is then 30 added to a 50  $\mu$ l amplification reaction as described in Example 2. The resulting



WO 00/41524

PCT/US00/00580

31

amplified DNA is then sequenced as described in Example 4. While the centrifugation step removes genomic DNA and thus improves the purity of the sample, it is possible to avoid this step and use the lysed sample directly for amplification and sequencing.

While the lysis method described here uses lysozyme, other methods could be used, such as the addition of nonionic detergents (e.g. Triton X-100 or Tween 20) or the expression of the T7 lysozyme gene (gene 3.5 protein) within the *E. coli* cell (pLysS and pLysE, Novagen, Inc., Madison, WI). It is possible that the method of lysis will have to be optimized for plasmids of varying size in order to minimize the amount of contaminating chromosomal DNA released.

10

**Example 7: BAC end sequencing by selective amplification of the ends of the insert**

In this example the ends of BAC clones are sequenced by first preferentially amplifying DNA fragments containing the ends of each insert. This is accomplished by digesting the BAC DNA and then carrying out a ligation in the presence of a splint that will result in circularization of only the two end fragments. Since the amplification system described in Example 2 preferentially amplifies circular DNA when the DNA fragments are less than several kbp, this circularization results in an enrichment of the end fragments in the final amplified mixture. (This is not necessarily the case for other amplification systems of the invention.) While the example below uses the vector pBELO-11 (Research Genetics, Huntsville, AL), which has a single *Bae* I site and no *Mlu* I sites, the strategy used can readily be applied to other BAC vectors by choosing the appropriate restriction enzymes and designing the appropriate splint oligonucleotides.

pBELO-11 BAC DNA containing inserts is isolated from *E. coli* cells using standard procedures. 50 ng of the DNA is treated with the enzymes *Bae* I, *Mlu* I and T4 DNA ligase (New England Biolabs, Beverly, MA) and a 50-fold molar excess of the following three oligonucleotides: BAC-1 (5'-CGCGGTACACCGACGTCAA-3'), BAC-2 (5'-CGCGGTACACCGACTTAAT-3') and BAC-3 (5'-GTCGGTGTAC-3'). BAC-1 and BAC-3 will anneal to form a split that will result in circularization and ligation of one end of the pBELO-11 DNA, while BAC-2 and BAC-3 will anneal to form a split that will result in circularization and ligation of the other end of the pBELO-11 DNA. The

30

WO 00/41524

32

PCT/US00/00580

reactions are carried out in 20  $\mu$ l in 20 mM Tris-HCl, pH 7.5, 10 mM MgCl<sub>2</sub>, 5 mM DTT, 20  $\mu$ M S-adenosylmethionine and 500  $\mu$ M ATP at 37 °C for 60 min. 5  $\mu$ l of the ligated products are amplified as described in Example 2 in a 50  $\mu$ l reaction. After removal of the unincorporated nucleotides, the amplified DNA is sequenced using the universal T7 (TAATACGACTCACTATAGGGCGA) or SP6 (CATACGATTTAGGTGACACTATAG) primers that anneal upstream of each of the two ends of the insert in pBELO-11.

Example 8: Generic amplification of human genomic DNA

10 A cheek swab is taken of a human subject and applied to FTA Card (GIBCO BRL, Rockville, MD). The DNA bound to the paper is purified using FTA Purification Reagent as described by the distributor (GIBCO BRL, Rockville, MD). The DNA is eluted from the paper in 10 mM Tris-HCl, pH 7.5, 1 mM EDTA by heating at 80 °C for 5 min. The concentration of the eluted DNA is approximately 1 ng/ $\mu$ l. One  $\mu$ l of this  
15 DNA is amplified in a 50  $\mu$ l reaction as described in Example 2, incubating at 37 °C for 20 min. The concentration of the DNA in the completed reaction is 0.5  $\mu$ g/ $\mu$ l, a 25,000-fold amplification. Aliquots of this DNA can be used directly for genotyping using PCR reactions; e.g. 1  $\mu$ l in a 25  $\mu$ l PCR reaction.

20 Example 9: Detection of contaminating DNA by amplification reaction

In order to characterize the level of contamination of DNA in an unknown sample, reactions are carried out as described in Example 2 using varying amounts of the unknown sample. As a control, a standard series of reactions are also carried out both in the absence of DNA and in the presence of 1, 10, 100 and 1,000 pg of a standard  
25 supercoiled plasmid DNA such as pUC18. The reactions are carried out at 37°C for 15 min. The reaction mixtures are carried out in the presence of 20 cpm/pmol [3H]TTP, and the amount of DNA synthesis in each sample is determined by measuring the amount of radioactivity taken up into DNA, as described in Example 1. Alternatively, DNA synthesis can be monitored in real time using a fluorescent probe (see Example 4).  
30 The amounts of DNA synthesized in the unknown samples are compared with the

WO 00/41524

PCT/US00/00580

33

amounts of DNA synthesized in the standard reactions containing different initial concentrations of the known DNA in order to extrapolate the relative amount of contaminating DNA in the unknown sample.

5 Example 10: Single molecule amplification in agarose

In accordance with the description above, in this example, single DNA molecules are amplified as individual foci embedded in a solid matrix, such as agarose. The amplified DNA is detected by fluorescence in the presence of the dye SYBR Green II (Molecular Probes, Eugene, OR).

- 10 Reactions are carried out as described in Example 4. Two-fold concentrated reaction mixtures are prepared, and then diluted two-fold with a solution containing liquid 1% SeaPlaque agarose (FMC Products, Rockland, ME). The agarose is prepared in water and dissolved at 90 °C, then cooled to 37 °C before mixing with the reaction mixtures. The reaction mixtures contain a final 1/80,000 X concentration of SYBR
- 15 Green II fluorescent dye, and varying amounts of supercoiled pUC18 DNA, ranging from none up to 100,000 molecules per  $\mu$ l. After mixing the agarose and the reaction mixture, the mixture is plated on a microscope slide, covered with a cover slip, and placed on ice to solidify the mixture. The slide is then heated at 37 °C and the progression of foci is monitored using a fluorescence microscope, exciting at a
- 20 wavelength of 480 nM and observing the emission at 530 nM. In the absence of added DNA, approximately 20 foci are observed on a one  $\text{cm}^2$  region, that correspond to the amplification of contaminating DNA. In the presence of increasing amounts of pUC18 DNA, the number of foci increasing proportionally, and thus correspond to amplification of the added DNA. Each foci corresponds to the amplification of a single DNA
- 25 molecule, and thus is "clonal".

- While the example below uses supercoiled pUC18 DNA as a template, one skilled in the art will recognize that other circular DNAs could be used. For example, if BAC DNA was digested with EcoRI, and then ligated, a population of circular fragments, or a "library", would be generated. If a splint oligonucleotide was ligated
- 30 between the EcoRI sites, these could then serve as primer sites to sequence each insert in both directions. Using this strategy, all the amplified foci embedded on an agarose bed

WO 00/41524

PCT/US00/00580

34

could be sequenced in a single dye-terminator sequencing reaction. After carrying out the sequencing reactions, the unincorporated dye terminators and the salts could be washed out of the agarose and each foci could be sequenced directly by injection into a capillary-based sequencing instrument such as the PE Biosystems 3700. This would  
5 allow one to generate a library of subclones of a large DNA fragment rapidly without the use of *E. coli* cells, and then to sequence all the fragments in a single sequencing reaction, that could be directly loaded onto a capillary sequencing instrument. This would represent a tremendous reduction in the scale of required reagents and human labor involved in cloning and sequencing fragments.

10

Example 11: Overproduction and purification of the 63-kDa T7 gene 4 protein

A preferred T7 gene 4 protein used in this invention is the 63-kDa form referred to as G4A<sub>G64</sub> in Mendelman *et al.*, (*Proc. Natl. Acad. Sci. USA* 89:10638, 1992) and Mendelman *et al.*, (*J. Biol. Chem.* 268:27208, 1993). It is the wild-type 63-kDa gene 4  
15 protein except that the methionine at residue 64 has been replaced with a glycine to prevent initiation of synthesis of the 56-kDa form of the gene 4 protein. Expression of the T7 gene 4 in *E. coli* is toxic to the cells and must be kept tightly repressed when uninduced. Furthermore, when induced, the toxicity prevents efficient overproduction of the protein. It is known that mutants of gene 4 protein such as those in the nucleotide  
20 binding site eliminate this toxicity and result in much greater overproduction of the protein (Notarnicola and Richardson, *J. Biol. Chem.* 268:27198, 1993). This observation suggests that it is the potent TTPase activity of the wild-type gene 4 protein that is responsible for the toxicity. Thus one method to increase the amount of wild-type gene 4 protein overproduced in cells is to coexpress another molecule that will bind to the wild-  
25 type gene 4 protein and inhibit its TTPase activity. Three possible classes of molecules that may accomplish this are protein, RNA and DNA aptamers (Cohen *et al.*, *Proc. Natl. Acad. Sci. USA* 95:14272, 1998; Famulok and Jenne, *Curr. Opin. Chem. Biol.* 2:320, 1998), which can be selected for from libraries on the basis of either binding to wild-type gene 4 protein *in vitro*, or inhibit the toxicity of its production *in vivo*.

WO 00/41524

35

PCT/US00/00580

An alternative approach is to coexpress a fragment of gene 4 that interferes with the activity of the wild-type gene 4 protein. The 295 residue carboxyl-terminal fragment of gene 4 protein from arginine 271 to the end of the gene has this property; *in vitro* it inhibits the wild-type gene 4 protein's TTPase activity, and *in vivo* it reduces the toxicity of the wild-type gene 4 and increases the overproduction of the wild-type gene 4 protein by at least 10-fold (Guo *et al.*, J. Biol. Chem. 274:30303, 1999). It is likely that other carboxyl-terminal fragments of the gene 4 will have a similar effect.

To overproduce the G4A<sub>64A</sub> form of the gene 4 protein, the plasmid pGP4A/4E-1 whose sequence is shown in Figure 1 (SEQ. ID. NO. 1), was used. It expresses both the G4A<sub>64A</sub> gene 4 and the gene 4 fragment initiating at codon 271, both from T7 RNA polymerase promoters. The two genes are in tandem, with the G4A<sub>64A</sub> expressed upstream of the gene 4 fragment that starts at codon 271 (see Figure 1). pGP4A/4E-1 was used to transform the *E. coli* cells BL21/DE3 (Novagen, Inc., Madison, WI) and selected for by plating on agar plates containing 50 µg/ml kanamycin. The cells were grown in LB media containing 50 µg/ml kanamycin at 30°C. When the cell density reached A<sub>590</sub>=1, the gene 4 proteins were induced by the addition of 0.5 mM IPTG. After inducing for 3 hours at 30°C, the cells were harvested and frozen at -80°C.

The 63-kDa gene 4 protein was purified by the procedure described by Notarnicola *et al.* (J. Biol. Chem. 270:20215, 1995). Briefly, the cells were lysed by the addition of lysozyme and by sonication. After the cell debris was removed by centrifugation, the 63-kDa gene 4 protein was precipitated by the addition of 15% polyethylene glycol 4000. After resuspension of the precipitate, the gene 4 protein was purified by phosphocellulose chromatography. Finally, the pool of the fractions containing gene 4 protein from the phosphocellulose column was purified by ATP-agarose affinity chromatography. The gene 4 protein fragment present in the extract separates from the G4A<sub>64A</sub> gene 4 protein both in the PEG precipitation, the phosphocellulose chromatography, and the ATP affinity chromatography. From 10 liters of induced cell culture, 100 mg of pure G4A<sub>64A</sub> gene 4 protein were obtained.

Other embodiments within the scope of the following claims.

WO 00/41524

PCT/US00/00580

36

CLAIMS

What is claimed is:

1. A method of amplifying a template DNA molecule comprising:  
incubating said template DNA molecule with a reaction mixture  
5 comprising a DNA polymerase and at least one accessory protein at a constant  
temperature to produce amplified product, wherein production of amplified product  
does not require exogenously-added oligonucleotide primers and said template DNA  
molecule does not have a terminal protein covalently bound to either 5' end.
2. The method of claim 1, wherein said DNA polymerase is a  
10 bacteriophage DNA polymerase.
3. The method of claim 1, wherein said DNA polymerase is  
bacteriophage T7 DNA polymerase.
4. The method of claim 3, wherein said DNA polymerase is a mixture of  
wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced  
15 3' to 5' exonuclease activity.
5. The method of claim 1, wherein said accessory protein is a helicase.
6. The method of claim 1, wherein said accessory protein is a primase.
7. The method of claim 1, wherein said accessory protein is the  
helicase/primase of bacteriophage T7.
- 20 8. The method of claim 7, wherein said helicase/primase is the 63-kDa  
form of the protein from bacteriophage T7.
9. The method of claim 1, wherein said accessory protein is a single-  
stranded DNA binding protein.
- 25 10. The method of claim 9, wherein said single-stranded DNA binding  
protein is from *Escherichia coli*.

WO 00/41524

PCT/US00/00580

37

11. A method of amplifying a template DNA molecule comprising:  
incubating said template DNA molecule with an *in vitro* reaction  
mixture comprising a DNA polymerase, a helicase, and a primase at a constant  
temperature to produce amplified product.
- 5 12. The method of claim 11, wherein said reaction mixture does not  
require exogenously-added oligonucleotide primers.
13. The method of claim 11, wherein said polymerase is from  
bacteriophage T7.
14. The method of claim 13, wherein said DNA polymerase is a mixture of  
10 wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced  
3' to 5' exonuclease activity.
15. The method of claim 11, wherein said helicase and said primase is the  
helicase/primase of bacteriophage T7.
16. The method of claim 15, wherein said helicase/primase is the 63-kDa  
15 form of the protein from bacteriophage T7.
17. The method of claim 11, wherein said reaction mixture further  
comprises a single-stranded DNA binding protein.
18. The method of claim 17, wherein said single-stranded binding protein  
is from *Escherichia coli*.
- 20 19. The method of claim 11, where the amount of DNA produced is a  
1,000-fold increase over the amount of the template DNA.
20. The method of claim 11, where the amount of DNA produced is a  
10,000-fold increase over the amount of the template DNA.
21. The method of claim 11, where the amount of DNA produced is a  
25 100,000-fold increase over the amount of the template DNA.

WO 00/41524

PCT/US00/00580

38

22. The method of claim 11, where the amount of DNA produced is a 1,000,000-fold increase over the amount of the template DNA.

23. The method of claim 11, where the amount of DNA produced is a 10,000,000-fold increase over the amount of the template DNA.

5        24. A method of amplifying a template DNA molecule comprising:  
incubating said template DNA molecule in an *in vitro* reaction mixture  
comprising a wild-type T7 DNA polymerase and a T7 DNA polymerase modified to  
have reduced 3' to 5' exonuclease activity, a 63-kDa form of a gene 4 protein from  
bacteriophage T7 and a single-stranded binding protein from *Escherichia coli* at a  
10       constant temperature to produce amplified product.

25. The method of claim 24, wherein said reaction mixture does not  
require exogenously-added oligonucleotide primers.

26. The method of claims 1, 11, or 24, wherein said amplification of said  
template DNA is exponential.

15       27. The method of claims 1, 11, or 24, wherein said reaction mixture  
further comprises a single-stranded DNA binding protein of bacteriophage T7.

28. The method of claims 1, 11 or 24, wherein said reaction mixture  
further comprises a nucleoside diphosphokinase.

29. The method of claim 28, wherein said nucleoside diphosphokinase is  
20       from *Escherichia coli*.

30. The method of claims 1, 11 or 24, wherein said reaction mixture  
further comprises an inorganic pyrophosphatase.

31. The method of claims 1, 11 or 24, wherein said reaction mixture  
further comprises an ATP regeneration system.



WO 00/41524

PCT/US00/00580

39

32. The method of claim 31, wherein said ATP regeneration system consists of phosphocreatine and creatine kinase.

33. The method of claims 1, 11 or 24, wherein said reaction mixture further comprises a 5' to 3' exonuclease.

5 34. The method of claim 33, wherein said exonuclease is T7 gene 6 exonuclease.

35. The method of claims 1, 11 or 24, wherein said reaction mixture further comprises a DNA ligase.

36. The method of claim 35, wherein said ligase is T7 DNA ligase.

10 37. The method of claims 1, 11 or 24, wherein said template DNA molecule has a length of at least 1,000 bp.

38. The method of claims 1, 11 or 24, wherein said template DNA molecule has a length of at least 1,500 bp.

15 39. The method of claims 1, 11 or 24, where in said template DNA molecule has a length of at least 2,000 bp.

40. The method of claims 1, 11 or 24, wherein said template DNA molecule has a length of at least 3,000 bp.

41. The method of claims 1, 11 or 24, wherein said template DNA molecule is amplified at least about 10-fold.

20 42. The method of claims 1, 11 or 24, wherein said template DNA molecule is amplified at least about 100-fold.

43. The method of claims 1, 11 or 24, wherein said template DNA molecule is amplified at least about 1000-fold.

WO 00/41524

40

PCT/US00/00580

44. The method of claims 1, 11 or 24, wherein said constant temperature is less than 60° C.

45. The method of claims 1, 11 or 24, wherein said constant temperature of less than 45° C.

5 46. The method of claims 1, 11 or 24, wherein said amplification requires the presence of ATP and CTP.

47. The method of claims 1, 11 or 24, wherein said reaction mixture further comprises 0.5 to 5% DMSO.

48. The method of claims 1, 11 or 24, wherein said reaction mixture  
10 contains 0.5-5% dextran.

49. The method of claims 1, 11 or 24, wherein said template DNA molecule is from lysed *E. coli* cells.

50. The method of claim 49, wherein said DNA is a plasmid.

51. The method of claim 49, wherein said DNA is a bacterial artificial  
15 chromosome (BAC).

52. The method of claim 1, further comprising the step of treating said polymerase and at least one accessory protein with ultraviolet light prior to incubating with said template DNA molecule.

53. The method of claim 11, further comprising the step of treating said  
20 polymerases, said helicase, and said primase with ultraviolet light prior to incubating with said template DNA molecule.

54. The method of claim 24, further comprising the step of treating said polymerases, said gene 4 protein and said single-stranded binding protein with ultraviolet light prior to incubating with said template DNA molecule.

WO 00/41524

PCT/US00/00580

41

55. The method of claims 52, 53, or 54, wherein said ultraviolet light is a dose from 10 to 1000  $\mu\text{W}/\text{cm}^2$  for from 15 sec to 5 min.

56. The method of claim 1, 11 or 24, wherein said template DNA is less than 100 ng of a plasmid and said reaction mixture is 10 to 200  $\mu\text{l}$ .

5 57. A method for preparing a DNA molecule for use in a DNA sequencing reaction comprising the steps of:

incubating said DNA molecule in an *in vitro* reaction mixture comprising a DNA polymerase, a helicase, and a primase at a constant temperature so as to produce amplified product; and

10 providing said amplified product for use in a DNA sequencing reaction.

58. The method of claim 57, wherein said reaction mixture further comprises a single-stranded binding protein.

59. The method of claim 57, wherein said reaction mixture does not  
15 require exogenously-added oligonucleotide primers.

60. The method of claim 57, wherein production of said amplified product is exponential.

61. The method of claim 57, wherein the amount of said amplified product is at least 10-fold greater than said DNA molecule put into said reaction mixture.

20 62. A method for preparing a DNA molecule for use in a DNA sequencing reaction comprising the steps of:

incubating said DNA molecule in an *in vitro* reaction mixture comprising a T7 polymerase and a T7 gene 4 protein at a constant temperature so as to produce amplified product; and

25 providing said amplified product for use in a DNA sequencing reaction.

WO 00/41524

PCT/US00/00580

42

63. The method of claim 62, wherein said polymerase is a mixture of wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced 3' to 5' exonuclease activity.

64. The method of claim 62, wherein said T7 gene 4 protein is 63-kDa  
5 form of the protein.

65. The method of claim 62, wherein said reaction mixture further comprises a single-stranded binding protein from *Escherichia coli*.

66. The method of claim 62, wherein the amount of said amplified product is amplified is at least 10-fold greater than said DNA molecule put into said reaction  
10 mixture.

67. The method of claims 57 or 62, wherein said production of said amplified product is exponential.

68. The method of claim 62, wherein said reaction does not require exogenously-added oligonucleotide primers

69. The method of claims 57 or 62, wherein said the DNA sequencing  
15 reaction uses fluorescent primers.

70. The method of claims 57 or 62, wherein said DNA sequencing reaction uses fluorescent dideoxynucleotides.

71. The method of claims 57 or 62, wherein dNTPs present in said reaction  
20 mixture are removed after amplification by treatment with phosphatase.

72. The method of claim 71, wherein said phosphatase is from arctic shrimp.

73. The method of claims 57 or 62, wherein said DNA molecule is from lysed *E. coli* cells.

WO 00/41524

PCT/US00/00580

43

74. The method of claim 73, wherein said DNA is a plasmid.
75. The method of claim 73, wherein said DNA is a bacterial artificial chromosome (BAC).
76. The method of claim 73, wherein said DNA is from a human patient.
- 5 77. The method of claims 57 or 62, wherein said reaction mixture further comprises DMSO at a concentration of 0.5 to 8%.
78. The method of claims 57 or 62, wherein said reaction mixture further comprises dextran at a concentration of between 0.5 and 20%.
- 10 79. A kit for amplification, comprising:  
a DNA polymerase;  
a primase;  
a helicase; and  
a single-stranded binding protein.
80. The kit of claim 79, wherein said polymerase is T7 DNA polymerase.
- 15 81. The kit of claim 79, wherein said primase and helicase are a gene 4 protein from bacteriophage T7.
82. The kit of claim 79, wherein said single-stranded binding protein is from *Escherichia coli*.
- 20 83. The kit of claim 80, wherein said T7 polymerase is a mixture of a wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced 3' to 5' exonuclease activity.
84. The kit of claim 81, wherein said gene 4 protein is the 63-kDa form of the protein.
85. The kit of claim 84, further comprising a nucleoside diphosphokinase.

WO 00/41524

44

PCT/US00/00580

86. The kit of claim 85, wherein said nucleoside diphosphokinase is from *Escherichia coli*.
87. The kit of claim 79, further comprising inorganic pyrophosphatase.
88. The kit of claim 79, further comprises an ATP regeneration system.
- 5 89. The kit of claim 88, wherein said ATP regeneration system consists of phosphocreatine and creatine kinase.
90. The kit of claim 79, further comprising a T7 gene 6 exonuclease.
91. The kit of claim 79, further comprising a T7 DNA ligase.
92. The kit of claim 79, further comprising a single-stranded DNA binding  
10 protein of bacteriophage T7.
93. Method for overproducing the gene 4 protein of bacteriophage T7 comprising:  
coproducing the gene 4 protein in *E. coli* with an agent that increases its level of production.
- 15 94. The method of claim 93, wherein said agent is a carboxy terminal fragment of gene 4 protein.
95. The method of claim 94, wherein said carboxy terminal fragment of gene 4 comprises the carboxy terminus of the protein after residue 221 (glutamine).
96. The method of claim 95, wherein said carboxy terminal fragment of  
20 gene 4 comprises the carboxy terminus of the protein between residues 260 (tryptophan) and residue 280 (serine).
97. Method of determining the amount of contaminating DNA in a sample comprising the steps of:

WO 00/41524

PCT/US00/00580

45

incubating said sample potentially containing contaminating DNA with a reaction mixture comprising a DNA polymerase such that if said contaminating DNA is present said DNA molecule is amplified to produce amplified product;

incubating at least one pre-determined amount of control DNA with  
5 said reaction mixture comprising a DNA polymerase such that said control DNA molecule is amplified to produce amplified control product; and

comparing the amount of amplified product in said sample with the amount of amplified control product as an indication of the amount of contaminating DNA in said sample.

10 98. The method of claim 97, wherein said DNA polymerase is from bacteriophage T7.

99. The method of claim 97, wherein said reaction mixture is at constant temperature.

100. The method of claim 97, wherein said reaction mixture further  
15 comprises a helicase, a primase and a single-stranded binding protein.

101. The method of claim 97, wherein said control DNA is amplified at least 10-fold.

102. The method of claim 97, wherein said reaction does not require exogenously-added oligonucleotide primers.

20 103. Method of claim 97, wherein amplification of said control DNA is exponential.

104. Method for amplifying DNA directly from cells, comprising the steps of:

lysing said cells to form a lysate; and  
25 incubating said lysate in a reaction mixture comprising a DNA polymerase, a helicase, and a primase at a constant temperature, such that DNA present in said lysate is amplified to produce amplified product, wherein production

WO 00/41524

PCT/US00/00580

46

of amplified DNA product does not require exogenously-added oligonucleotide primers.

105. The method of claim 104, wherein said cells are bacterial cells.

106. The method of claim 104, wherein said cells are human cells.

5 107. The method of claim 104, wherein said reaction mixture further comprises a single-stranded binding protein.

108. Method for amplifying DNA directly from cells comprising the steps of:

lysing said cells to form a lysate; and

10 incubating said lysate in a reaction mixture comprising a DNA polymerase such that DNA present in said lysate is amplified to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers, said amplification is exponential and wherein the amount of amplified product is at least ten-fold greater than the amount of DNA present in said  
15 lysate put into said reaction mixture.

109. The method of claim 108, wherein said cells are bacterial cells.

110. The method of claim 108, wherein said cells are human cells.

111. The method of claim 49, wherein said DNA is from bacteriophage.

112. A method of amplifying a plasmid template DNA molecule,  
20 comprising incubating said template DNA molecule in a reaction mixture comprising a DNA polymerase to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers, is exponential, and the amount of amplified product is at least 10-fold greater than the amount of template DNA put into the reaction mixture.



WO 00/41524

PCT/US00/00580

47

113. A method for preparing a DNA molecule for use in a DNA sequencing reaction comprising the steps of:

- incubating said DNA molecule in a reaction mixture comprising a DNA polymerase to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers, is exponential, the amount of amplified product is at least 10-fold greater than the amount of DNA molecule put into the reaction mixture; and
- providing said amplified product for use in a DNA sequencing reaction.

- 10 114. The method of claim 108, wherein said amount of said amplified product

is at least 100-fold greater than the amount of DNA present in said lysate added to said reaction mixture.

- 15 115. The method of claim 108, wherein said amount of amplified product is at

least 1000-fold greater than the amount of DNA present in said lysate added to said reaction mixture.

116. The method of claim 112, wherein said amount of amplified product is at

- 20 least 100-fold greater than the amount of template DNA added to said reaction mixture.

117. The method of claim 112, wherein said amount of amplified product is at

- 25 least 1000-fold greater than the amount of template DNA present added to said reaction mixture.

WO 00/41524

48

PCT/US00/00580

118. The method of claim 113, wherein said amount of amplified product is at

least 100-fold greater than the amount of DNA added to said reaction mixture.

119. The method of claim 113, wherein said amount of amplified product is at

least 1000-fold greater than the amount of DNA added to said reaction mixture.

120. The method of claim 96, wherein said carboxy terminal fragment of gene 4 initiates at residue 271 (arginine).

121. A method for sequencing a nucleic acid sequence, wherein the DNA molecule to be sequenced is prepared in a constant temperature amplification reaction comprising the steps of:

incubating a single DNA molecule to be sequenced in a reaction mixture comprising a DNA polymerase to produce amplified product, wherein production of amplified product is done isothermally, is exponential, the amount of amplified product is at least 10-fold greater than the amount of DNA molecule put into the reaction mixture.

122. A method for sequencing wherein the DNA molecule to be sequenced is prepared in a reaction comprising the steps of:

incubating a single DNA molecule to be sequenced in a reaction mixture comprising a DNA polymerase to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers, is exponential, the amount of amplified product is at least 10-fold greater than the amount of DNA molecule put into the reaction mixture.

123. The method of claim 121, wherein said method of DNA preparing a DNA molecule for sequencing is performed isothermally.

WO 00/41524

PCT/US00/00580

1/3

Figure 1.

	10	20	30	40	50	60	70	80
ATGGACAATT	CGCAGGATTC	CGATAGTGTA	TTTCTTTACC	ACATTCCTTG	TGACAACTGT	GGGAGTAGTG	ATGGGAAGTC	80
GCTGTTCTCT	GACGGACACA	CGTTCTGCTA	CGTATGCGAG	AAGTGGACTG	CTGGTAATGA	AGACACTAAA	GAGAGGGCTT	160
CAAAACGGAA	ACCCTCCGGC	GGAAAGCCCG	GGACTTACAA	CGTGTGGAAC	TTCCGGGAAT	CCAATGGACG	CTACTCCGCG	240
TTAAGTCGA	GAGGAATCTC	CAAGGAAACC	TGTCAGAAAG	CTGGCTACTG	GATTGCCAAA	GTAGACGGTG	TGATGTACCA	320
AGTGGCTGAC	TATCGGGACC	AGAACGGCAA	CATTGTGAGT	CAGAAGGTTT	GAGATAAAGA	TAAGAACTTT	AAGACCACTG	400
GTAGTCACAA	CAGTGACGCT	CTGTTCCGGA	AGCACTTGTC	GAATGGTGGT	AAGAAGATTG	TCGTTACAGA	AGGTGAAATC	480
GACATGCTTA	CCGTGATGGA	ACTTCAAGAC	TGTAAGTATC	CTGTAGTGTG	GTGGGTGAC	GGTGCCTCTG	CCGTAAAGAA	560
GACATGCGCT	GCCAACTACG	AATACTTTGA	CCAGTTCGAA	CAGATTATCT	TAATGTTTGA	TATGGACGAA	GCAGGGCGCA	640
AAGCAGTCGA	AGAGGCTGCA	CAGGTTCAC	CTGCTGGTAA	GGTACGAGTG	GCAGTTCCTC	CGTGAAGGA	TGCAACGAG	720
TGTCACCTAA	ATGGTCACGA	CCGTGAAATC	ATGGAGCAA	TGTGGAATGC	TGGTCCCTGG	ATTCTGTATG	GTGTGGTATC	800
GGCTCTTTCC	TTACGTGAAC	GAATCCGTGA	GCACCTATCG	TCCGAGGAAT	CAGTAGGTTT	ACTTTTCAGT	GGCTGCACCTG	880
GTATCAACGA	TAAGACCTTA	GGTCCCCGTG	GTGGTGAAGT	CATTATGGTC	ACTTCCGGTT	CCGGTATGGG	TAAGTCAACG	960
TTGCTCCGTC	AACAAGCTCT	ACAATGGGGC	ACAGCGATGG	GCAAGAAGGT	AGGCTTAGCG	ATGCTTGAGG	AGTCCGTGGA	1040
GGAGACCGCT	GAGGACCTTA	TAGGTCTACA	CAACCGTGTC	CGACTGAGAC	AATCCGACTC	ACTAAACAGA	GAGATTATTG	1120
AGAACGGTAA	GTTCGACCAA	TGGTTCGATG	AACGTGTCGG	CAACGATACG	TTCCATCTAT	ATGACTCATT	CGCCGAGGCT	1200
GAGACGGATA	CACTGCTCGC	TAAGCTGGCC	TACATGCGCT	CAGGCTTGGG	CTGTGACGTA	ATCATTCTAG	ACCACATCTC	1280
AATCGTCGTA	TCCGCTTCTG	GTGAATCCGA	TGAGCGTAAG	ATGATTGACA	ACCTGATGAC	CAAGCTCAAA	GGGTTCCGTA	1360
AGTCAACTGG	GGTGGTGCTG	GTGTAATTTT	GTACCTTAA	GAACCCAGAC	AAAGGTAAAG	CACATGAGGA	AGGTCCGCCC	1440
GTTCCTATTA	CTGACCTACG	TGGTCTGCGC	GCACCTACGC	AACATCTGTA	TACTATTATT	GCCCTTGAGC	GTAATCAGCA	1520
AGGCGATATG	CCTAACCTTG	TCCTCGTTCC	TATCTCAAG	TGCCGCTTTA	CTGGTGATAC	TGGTATCGCT	GGCTACATCG	1600
AATACAACAA	GGAAACCGGA	TGGCTTGAAC	CATCAAGTTA	CTCAGGGGAA	GAAGACTCAC	ACTCAGAGTC	AACGAGCTGG	1680
TCCAACGACA	CTGACTTCTG	ACAGGATTCT	TGATGACTTT	CCAGACGACT	ACGAGAAGTT	TGCTGGGAGA	GTCCCATTTCT	1760
AATACGACTC	ACTAAAGGAG	ACACACCTGA	TTCAAACCTA	TTAAGAAGTT	AGGCCAACTG	CTGGTTCGTA	TGTACAACGT	1840
CGAAGCCCAAG	CGACTGAACG	ATGAGGCTCG	TAAAGAGGCC	ACACAGTCAC	GGCTCTGGCC	GATTGCTGCC	AAAAGTGGTT	1920
TTGCGCTTAC	CCCAACCAAC	AGGGGATTTG	CTGCTTTCCA	TTGACCTGT	TTCTCTGGCC	GACGTTCCGG	CGGGCGTGTT	2000
TGTGCATCCA	TCTGGATTCT	CCTGTCAGTT	AGCTTTGGTG	GTGTGTGGCA	GTGTAGTCC	TGAACGAAAA	CCCCCGCGA	2080
TTGGCACATT	GGCAGCTAAT	CCGGAATCGC	ACTTACGGCC	AATGCTTCGT	TTCTATCAC	ACACCCCAAA	GCCTTCTGCT	2160
TTGAATGCTG	CCCTTCTTCA	GGGCTTAATT	TTTAAGAGCG	TCACCTTCAT	GGTGGTCAGT	GGCTCCTGCT	GATGTGCTCA	2240
GTATCACCGC	CAGTGGTATT	TATGTCAACA	CCGCCAGAGA	TAATTTATCA	CCGCAGATGG	TTATCTGTAT	GTTTTTTATA	2320
TGAATTTATT	TTTTGACGGG	GGGCATTGTT	TGGTAGGTGA	GAGATCCGGC	TGCTAACAAA	GGCCGAAAGG	AAGCTGAGTT	2400
GGCTGCTGCC	ACCGCTGAGC	AATAACTAGC	ATAACCCCTT	GGGGCTCTA	AACGGGTCTT	GAGGGGTTTT	TTGCTGAAAG	2480
GAGGAAGTAT	ATCCGGATAT	CCCGCAAGAG	GCCTGGCAGT	ACCGGCATAA	CCAAGCTAT	GCCTACAGCA	TCCAGGGTGA	2560
CGGTGCCGAG	GATGACGATG	AGCGCATTTG	TAGATTTTAT	ACACGGTGCC	TGACTGCGTT	AGCAATTTAA	CTGTGATAAA	2640
CTACCGCATT	AAAGCTTCCG	CCCGCACTCC	ACGAACCCCTT	CGGATCTCGA	TCCCGCGAAA	TTAATACGAC	TCACTATAGG	2720
GAGACCAACA	CGGTTTCCCT	CTAGAATAAA	TTTTGTTTAA	CTTTAAGAA	GAGATATACA	TATGCGTGAA	CGAATCCGTG	2800
AGCACCTATC	GTCCGAGGAA	TCAGTAGGTT	TACTTTTCAG	TGGCTGCACT	GGTATCAACG	ATAAGACCTT	AGGTGCCCGT	2880
GGTGGTGAAG	TCATTATGGT	CACTTCCGGT	TCCGGTATGG	GTAAGTCAAC	GTTCTCCGT	CAACAAGCTC	TACAATGGGG	2960
CACAGCGATG	GGCAAGAAGG	TAGGCTTAGC	GATGCTTGAG	GAGTCCGTTG	AGGAGACCGC	TGAGGACCTT	ATAGGCTTAC	3040
ACAAACCGTGT	CCGACTGAGA	CAATCCGACT	CACTAAAGAG	AGAGATTATT	GAAGAACGTA	AGTTCCGACCA	ATGGTTTCGAT	3120
GAAGTGTTCG	GCAACGATAC	GTTCCATCTA	TATGACTCAT	TGCGCGAGGC	TGAGACGGAT	AGACTGCTCG	CTAAGCTGGC	3200
CTACATGCGC	TCAGGCTTGG	GCTGTGACGT	AATCACTCTA	GACCACATCT	CAATCGTCTG	ATCCGCTTCT	GGTGAATCCG	3280
ATGAGCGTAA	GATGATTGAC	AACCTGATGA	CCAAGCTCAA	AGGGTTCGCT	AAGTCAACTG	GGGTGGTCTG	GGTCTGTAAT	3360
TGTCACCTTA	AGAACCCAGA	CAAGGTAA	GCACATGAGG	AAGGTCGCCC	CGTTTCTATT	ACTGACCTAC	GTGGTCTCTG	3440
CGCACTACGC	CAACTATCTG	ATACTATTAT	TGCCCTTGAG	CGTAATCAGC	AAGCGGATAT	GCCTAACCTT	GTCTCTGTTT	3520

WO 00/41524

PCT/US00/00580

2/3

```

GTATTCTCAA GTGCCGCTTT ACTGGTGATA CTGGTATCGC TGGCTACATG GAATACAACA AGGAAACCGG ATGGCTTGAA 3600
CCATCAAGTT ACTCAGGGGA AGAAGAGTCA CACTCAGAGT CAACAGACTG GTCCAACGAC ACTGACTTCT GAGGATCCAC 3680
TAGTAACGGC CGCCAGTGTG CTGGAATTCT GCAGATATCC ATCACACTGG CGGCCGCTCG AGCACCACCA CCACCACCAC 3760
TGAGATCCGG CTGCTAACAA AGCCCGAAAG GAAGCTGAGT TGGCTGCTGC CACCGCTGAG CAATAACTAG CATAACCCCT 3840
TGGGGCCTCT AAACGGGTCT TGAGGGGTTT TTGCTGAAA GGAGGAAC TAATCCGGATT GCGGAATGGG ACGCCGCCCTG 3920
TAGCGGCGCA TTAAGCGCGG CGGGTGTGGT GGTACGCGC AGCGTGACCG CTACACTTGC CAGCGCCCTA GCGCCGCTC 4000
CTTTCCGCTT CTTCCTTCC TTCTCGCCA CGTTCGCCG TTCTCCCGT CAAGCTCTAA ATCGGGGGCT CCTTTAGGG 4080
TTCCGATTTA GTGCTTTACG GCACCTCGAC CCCAAAAAAC TTGATTAGGG TGATGGTTCA CGTAGTGGG CATCGCCCTG 4160
ATAGACGGTT TTTCGCCCTT TGACGTTGGA GTCCACGTTT TTTAATAGTG GACTCTTGT CCAAACCTGA ACAACACTCA 4240
ACCCATATCT GGTCTATTCT TTGATTATAT AAGGGATTTT GCCGATTTC GCCTATTGGT TAAAAATGA GCTGATTAA 4320
CAAAAATTTA ACGCGAATT TAACAAAATA TTAACGTTTA CAATTTTCAG TGGCACTTT CGGGGAAATG TCGCGGAAAC 4400
CCCTATTGT TTAATTTTCT AAATACATTC AAATATGTAT CCGCTCATGA ATTAATCTT AGAAAACTC ATCGAGCATC 4480
AAATGAACT GCAATTTAT CATATCAGGA TTATCAATAC CATATTTTG AAAAAAGCGT TTCTGTAATG AAGGAGAAAA 4560
CTCAGCGAGG CAGTTCCTA GGATGGCAAG ATCCTGGTAT CGGTCTGCGA TTCCGACTCG TCCAACATCA ATACAACCTA 4640
TTAATTTCCC CTGCTCAAAA ATAAGGTTAT CAAGTGAGAA CATTACGCTC GTCATCAAAA TCACCTCGCAT CAACCAAAAC 4800
AGTTTATGCA TTCTTTTCCA GACTTGTTC ACAGGCCAGC CGATCGCTGT TAAAGGACA ATTACAACA GGAATCGAAT 4880
GTTATTCAAT CGTGATTGCG CCTGAGCGAG ACGAAATACG TTCACCTGAA TCAGGATATT CTCTAATAC CTGGAATGCT 4960
GCAACCGGCG CAGGAACACT GCCAGCGCAT CAACAATATT CAGGAGTACG GATAAAATGC TTGATGGTCG GAAGAGGCAT 5040
GTTTTCGCGG GGATCGCAGT GGTGAGTAAC CATGCATCAT ACATCATTTG CAACGCTACC TTTGCCATGT TTCAGAAACA 5120
AAATTCGCTC AGCCAGTTTA GTCTGACCAT CTCATCTGTA CGCACCTGAT TGCCCGACAT TATCGCGAGC CCATTTATAC 5200
ACTCTGGGCG ATCGGGCTTC CCATACAATC GATAGATTGT CCATATAAA' CAGCATCCAT GTTGGAA'IT AA'CGCGGCC 5280
CCTTGTATTA CTCTTTATCT AACAGACAG TTTTATGTG CATGACCAA ATCCCTTAA CTACTTTTC CTTCCTGTA 5360
GGGTGAGACC CCGTAGAAAA GATCAALGGA TCTTCTGAG ATCTTTTTT TCTGCGCGTA ATCTGCTGCT TGCAAAACA 5440
AAAAACCCG CTACCAAGCG GTGTTTGTTC GCCGATCAA GAGCTACCAA CTCTTTTTC GAAGGTAAT GGCTTCAGCA 5520
GAGCGCAGAT ACCAAATACT GTCTTCTAG TGTAGCCGTA GTTAGGCCAC CACTTCAAGA ACTCTGTAGC ACCGCTACA 5600
TACCTCGCTC TGCTAATCT GTTACCAAGT GCTGCTGCCA GTGGCGATAA GTCGTGCTT ACCGGGTGG ACTCAACAG 5680
ATAGTTACCG GATAAGGCGC AGCGGTGCGG CTGAAAGGGG GGTTCGTGCA CACAGCCAG CTGGAGCGA ACGACCTACA 5760
CCGAACGAG ATACCTACAG CGTGAGCTAT GAGAAAGCGC CACGCTTCCC GAAGGAGAA AGCGGACAG GTATCCGGA 5840
AGCGGACGGG TCGGAACAGC AGAGCGCAGC AGGGAGCTTC CAGGGGAAA CGCCTGTAT CTTTATAGTC CTGTCCGGTT 5920
TCGCCACCTC TGACTTGAGC GTGATTTT GTATGCTCG CTAGGGGGGG GGAGCCTATG GAAAAACGC AGCAACGCG 6000
CCTTTTACG GTTCTGGCC TTTTGTGCTA CATGTTCTT CTGCGTTAT CCCCTGATC TGTGGATAAC 6080
CGTATTACCG CCTTTGAGT AGCTGATACC GCTCGCCGCA CCGCAACGAC CGAGCGCAGC GAGTCAGTGA GCGAGGAAGC 6160
GGAAGAGCGC CTGATCGGT ATTTTCTCT TACGCATCTC TGCGGTATT CACACCGCAT ATATGGTGCA CTCTCAGTAC 6240
AATCTGCTCT GATGCCGAT AGTTAAGCCA GTATACATCT CCGTATCGCT ACGTGACTG GTCATGGCTG GCGCCGACA 6320
CCCGCCAACA CCGCTGACG CGCCCTGACG GGTGTGCTG CTCCCGCAT CCGCTTACG ACAAGCTGTG ACCGTCTCG 6400
GGAGCTGCAT GTGTCAGAG TTTTCAACGT CATCACCGAA ACGCGGAGG CAGCTCGGT AAAGCTCATC AGCGTGGTCG 6480
TGAAGCGATT CACAGATGTC TGCTGTTC TCCGCTCCA GCTCGTTGAG TTTCTCCAGA AGCGTTAATG TCTGGCTTCT 6560
GATAAGCGG GCCATGTTAA GGGCGGTTT TTCTGTTC GTCACTGAT CCTCCGTGA AGGGGATTT CTGTTTATG 6640
TGTGAGGGTA AACAACCTGG GGTATGGATG CGCGGGGACC ACGGTTACT GATGATGAAC ATGCCCGGT ACTGGAACGT 6720
TACAGATGTA GGTGTTCCAC AGGTAGCCA GCAGCATCTT AGAGAAAAAT CACTCAGGGT CAATGCCAGC GCTTCGTTAA 6800
TCCGCGTTTC CAGACTTTAC GAAACACGGA AACCGAAGAC CCGATGAGA TCCGGAACAT AATGGTGAG GCGCTGACT 6880
CAGTCTCTC ACGTTCGCT CCGTATCGGT GATTCATTCT GCTAACCAGT AAGGCAACCC CGCCAGCCTA GCGGGTCTT 7040
CAACGACAGC AGCAGATCA TGCGCACCCG TGGGGCCGCC ATGCGCGGCA TAATGGCCTG CTCTCGCCG AACGTTTGG 7120
TGGCGGGACC AGTGACGAAG GCTTGAGCGA GGGCGTGCAA GATTCCGAAT ACCGCAAGCG ACAGGCGAT CATGCTCGC 7200
CTCCAGCGAA AGCGGTCTC GCGCAAAATG ACCCAAGCG AGTCCGCGAC CTGCTCTACG AGTTGCATGA TAAAGAGAC 7280
AGTCATAAGT GCGGCGACGA TAGTCATGCC CCGCGCCAC CGGAAGGAGC TGACTGGGT GAAGGCTCTC AAGGGCATCG 7360
CTCGAGATCC CGGTGCTTAA TGAGTGAGCT AACTTACAT AATTGCGTTG CGCTCACTGC CCGCTTTCCA GTCGGGAAAC 7440
GTGTCTGTC AGCTGCATTA ATGAATCGGC CAACGCGCG GGAGAGGCGG TTTGCGTATT GGGCGCCAG GTGGTTTTTC 7520
TTTTACCAAG TGAGACGGGC AACAGCTGAT TGCCCTTCAC CGCTTGGCCC TGAGAGAGTT GCAGCAAGCG GTCCACGCTG 7600
GTTTGCCCA GCAAGCGGAA ATCTGTTTG ATGGTGGTTA ACGGCGGAT ATAACATGAG CTGCTCTCG TATCGTCGTA 7680
TCCCACTACC GAGATATCC CACCAACGCG CAGCCCGGAC TCGGTAAAG GCGCATTCG GCCCAGCGC ATCTGATCGT 7760
TGGCAACCA GATCGCAGTG GGAACGATG CCTCATTCAG CATTTGCATG GTTTGTTGAA AACCGACAT GGCCTCCAG 7840

```

WO 00/41524

PCT/US00/00580

3/3

```
TCGCCTTCCC GTTCCGCTAT CGGCTGAATT TGATTGCGAG TGAGATATTT ATGCCAGCCA GCCAGACGCA GACGCGCCGA 7920
GACAGAACTT AATGGGCCCC CTAACAGCGC GATTTGCTGG TGACCCAATG CGACCAGATG CTCCACGCCC AGTCGCGTAC 8000
CGTCTTCATG GGAGAAAATA ATACTGTTGA TGGGTGCTCG GTCAGAGACA TCAAGAAATA ACGCGGAAC ATTAGTGCA 8050
GCAGCTTCCA CAGCAATGGC ATCCTGGTCA TCCAGCGGAT AGTTAATGAT CAGCCCACTG ACGCGTTGCG CGAGAAGATT 8160
GTGCACCGCC GCTTTACAGG CTTCCAGGCC GCTTCGTTCT ACCATCGACA CCACCACGCT GGCACCCAGT TGATCGGCGC 8240
GAGATTTAAT CGCCCGGACA ATTTGCGAGC GCGCGTGCGAG GGCCAGACTG GAGGTGGCAA CGCCAATCAG CAACGACTGT 8320
TTGCCCGCCA GTTGTGTTGC CACGCGGTTG GGAATGTAAT TCAGCTCCGC CATCGCCGCT TCCACTTTTT CCCGCGTTTT 8400
CGCAGAAACG TGGCTGGCCT GGTTCACCAC GCGGAAACG GTCTGATAAG AGACACCGGC ATACTCTGCG ACATCGTATA 8480
ACGTTACTGG TTTCACATTC ACCACCTGA ATTGACTCTC TTCGGGGCGC TATCATGCCA TACCGCGAAA GGTTTTGGCG 8560
CATTCGATGG TGTCCGGGAT CTCGACGCTC TCCCTTATGC GACTCCTGCA TTAGGAAGCA GCCCAGTAGT AGGTTGAGGC 8640
CGTTGAGCAC CGCCGCCGCA AGGAATGGTG CATGCAAGGA GATGGGCGCC AACAGTCCCC CGGCCAGGGG GCCTGCCACC 8720
ATACCCACGC CGAAACAAGC GCTCATGAGC CCGAAGTGGC GAGCCCGATC TTCCCATCG GTGATGTCGG CGATATAGGC 8800
GCCAGCAACC GCACCTGTGG CGCCGTTGAT GCCGGCCACG ATGCGTCCGG CGTAGAGGAT CGAGATCTCG ATCCCGCGAA 8880
ATTAATACGA CTCACTATAG GCGAATTGTG AGCGGATAAC AATTCCTCTC TAGAAATAAT TTTGTTTAA 8960
AGATATACAT 8970
```

WO 00/41524

PCT/US00/00580

1

## SEQUENCE LISTING

<110> TABOR, STANLEY  
RICHARDSON, CHARLES

<120> ISOTHERMAL AMPLIFICATION OF DNA

<130> 249/107 WO

<140> To Be Assigned  
<141> Filed Herewith

<150> 60/115,498  
<151> 1999-01-11

<160> 6

<170> FastSEQ for Windows Version 3.0

<210> 1  
<211> 8970  
<212> DNA  
<213> Bacteriophage

<400> 1

atgggacaatt	cgcacgattc	cgatagtgt	tttctttacc	acattccttg	tgacaactgt	60
gggagtagtg	atgggaactc	gctgttctct	gacggacaca	cgttctgcta	cgtagcgag	120
aagtggactg	ctggtaatga	agacactaaa	gagagggctt	caaaacggaa	accctccggt	180
ggaaaagccc	ggacttacaa	cgtgtggaac	ttcggggaat	ccaatggacg	ctactccgcg	240
ttaactgcga	gaggaatctc	caaggaaacc	tgtcagaagg	ctggctactg	gattgccaaa	300
gtagacggtg	tgatgtacca	agtggctgac	tatcgggacc	agaacggcaa	cattgtgagt	360
cagaagggtc	gagataaaga	taagaacttt	aagaccactg	gtagtcacaa	gagtgacgct	420
ctgttcggga	agcacttggt	gaatggtggt	aagaagattg	tcgttacaga	aggtgaaatc	480
gacatgctta	ccgtgatgga	acttcaagac	tgtaagtatc	ctgtagtgtc	gttgggtcac	540
ggtgcctctg	ccgctaagaa	gacatgcgct	gccaaactac	aatactttga	ccagttcgaa	600
cagattatct	taatgttcga	tatggacgaa	gcagggcgca	aagcagtcga	agaggctgca	660
caggttctac	ctgctggtaa	ggtacgagtg	gcagttcttc	cgtgtaagga	tgcaaacgag	720
tgtcacctaa	atggtcacga	ccgtgaaatc	atggagcaag	tgtggaatgc	tggtccttgg	780
attcctgatg	gtgtgggtatc	ggctctttcg	ttacgtgaac	gaatccgtga	gcacctatcg	840
tccgaggaat	cagtagggtt	acttttcagt	ggctgcactg	gtatcaacga	taagacctta	900
ggtgcccgtg	gtgtggaagt	cattatggtc	acttccggtt	ccggtatggg	taagtcaacg	960
ttcgtccgtc	aacaagctct	acaatggggc	acagcgatgg	gcaagaagg	aggcttagcg	1020
atgcttgagg	agtcggttga	ggagaccgct	gaggacctta	taggtctaca	caaccgtgtc	1080
cgactgagac	aatccgactc	actaaagaga	gagattattg	agaacggtaa	gttcgaccaa	1140
tggttcgatg	aactgttcgg	caacgatacg	ttccatctat	atgactcatt	cgccgaggct	1200
gagacggata	gactgctcgc	taagctggcc	tacatgcgct	caggcttggg	ctgtgacgta	1260

WO 00/41524

PCT/US00/00580

2

atcattctag	accacatctc	aatcgctcga	tccgcttctg	gtgaatccga	tgagcgtaag	1320
atgattgaca	acctgatgac	caagctcaaa	gggttcgcta	agtcactgg	ggtggtgctg	1380
gtcgtaattt	gtcaccttaa	gaaccagac	aaaggtaaag	cacatgagga	aggtcgcccc	1440
gtttctatta	ctgacctacg	tggttctggc	gcactacgcc	aactatctga	tactattatt	1500
gcccttgagc	gtaatcagca	aggcgatatg	cctaaccttg	tcctcgttcg	tattctcaag	1560
tgccgcttta	ctggtgatac	tggtatcgct	ggctacatgg	aatacaacaa	ggaaaccgga	1620
tggttggaac	catcaagtta	ctcaggggaa	gaagagtcac	actcagagtc	aacagactgg	1680
tccaacgaca	ctgacttctg	acaggattct	tgatgacttt	ccagacgact	acgagaagtt	1740
tcgctggaga	gtcccattct	aatacgactc	actaaaggag	acacaccatg	ttcaaactga	1800
ttaagaagtt	aggccaactg	ctggttcgta	tgtacaacgt	ggaagccaag	cgactgaacg	1860
atgaggctcg	taaagaggcc	acacagtcac	gcgctctggc	gattcgctcc	aaaactgggt	1920
ttgcgcttac	cccaaccaac	aggggatttg	ctgctttcca	ttgagcctgt	ttctctgcgc	1980
gacgttcgcg	ccgcgctggt	tgtgcattcc	tctggattct	cctgtcagtt	agctttggtg	2040
gtgtgtggca	gtttagtacc	tgaacgaaaa	ccccccgcga	ttggcacatt	ggcagctaata	2100
ccggaatcgc	acttacggcc	aatgcttcgt	ttcgatatcac	acaccccaaa	gccttctgct	2160
ttgaatgctg	cccttcttca	gggcttaatt	tttaagagcg	tcaccttcat	ggtggtcagt	2220
gcgtcctgct	gagtgtctca	gtatcacccg	cagtggtatt	tatgtcaaca	ccgcagaga	2280
taatttatca	ccgcagatgg	ttatctgtat	gttttttata	tgaatttatt	ttttgcaggg	2340
gggcattggt	tggtagggtg	gagatccggc	tgctaacaaa	gcccgaagg	aagctgagtt	2400
ggctgctgcc	accgctgagc	aataactagc	ataacccctt	ggggcctcta	aacgggtcct	2460
gaggggtttt	ttgctgaaag	gaggaactat	atccggatat	cccgaagag	gcccggcagt	2520
accggcataa	ccaagcctat	gcctacagca	tccagggtga	cggtgccgag	gatgacgatg	2580
agcgacttgt	tagatttcat	acacggtgcc	tgactgcgtt	agcaatttaa	ctgtgataaa	2640
ctaccgcatt	aaagcttcgc	gcccgaactc	acgaaccctt	cggatctcga	tcccgcgaaa	2700
ttaatacgac	tcactatagg	gagaccacaa	cggtttccct	ctagaaataa	ttttgtttaa	2760
ctttaagaag	gagatataca	tatgcgtgaa	cgaatccgtg	agcacctatc	gtccgaggaa	2820
tcagtaggtt	tacttttctg	tggtgcact	ggatcaacg	ataagacctt	aggtgcccg	2880
ggtggtgaag	tactattggt	cacttccggt	tccggtatgg	gtaagtcaac	gttcgtccgt	2940
caacaagctc	tacaatgggg	cacagcgatg	ggcaagaagg	taggcttagc	gatgcttgag	3000
gagtcggttg	aggagaccgc	tgaggacctt	ataggtctac	acaaccgtgt	ccgactgaga	3060
caatccgact	cactaaagag	agagattatt	gagaacggtg	agttcgacca	atggttcgat	3120
gaactgttcg	gcaacgatac	gttccatcta	tatgactcat	tcgccgaggg	tgagacggat	3180
agactgctcg	ctaagctggc	ctacatgcgc	tcaggcttgg	gctgtgacgt	aatcattcta	3240
gaccacatct	caatcgctgt	atccgcttct	gggtgaatccg	atgagcgtaa	gatgattgac	3300
aacctgatga	ccaagctcaa	agggttcgct	aagtcaactg	gggtggtgct	ggtcgtaatt	3360
tgtcacctta	agaaccaga	caaaggtaaa	gcacatgagg	aaggtcgccc	cgtttctatt	3420
actgacctac	gtggttctgg	cgcactacgc	caactatctg	atactattat	tgcccttgag	3480
cgtaatcagc	aaggcgatat	gcctaaccct	gtcctcgctc	gtattctcaa	gtgccgcttt	3540
actggtgata	ctggtatcgc	tggtacatg	gaatacaaca	aggaaaccgg	atggcttgaa	3600
ccatcaagtt	actcagggga	agaagagtca	cactcagagt	caacagactg	gtccaacgac	3660
actgacttct	gaggatccac	tagtaacggc	cgccagtgtg	ctggaattct	gcagatatcc	3720
atcacactgg	cggccgctcg	agcaccacca	ccaccaccac	tgagatccgg	ctgctaacaa	3780
agcccgaag	gaagctgagt	tggtcgtgct	caccgctgag	caataactag	cataaccctt	3840
tggggcctct	aaacgggtct	tgaggggttt	tttgcgtgaa	ggaggaacta	tatccggatt	3900
ggcgaaatggg	acgcgccctg	tagcggcgca	tttagcgcg	cggtgtggtg	ggttacgcgc	3960
agcgtgaccg	ctacacttgc	cagcgcccta	gcgccgctc	ctttcgcttt	cttcccttcc	4020
tttctcgcca	cgttcgccgg	ctttccccgt	caagctctaa	atcggggggt	ccctttaggg	4080
ttccgattta	gtgcttttac	gcacctcgac	cccaaaaaac	ttgattaggg	tgatggttca	4140
cgtagtgggc	catcgccctg	atagacgggt	tttcgccctt	tgacgttgga	gtccacgttc	4200
tttaattagtg	gactcttgtt	ccaaactgga	acaacactca	accctatctc	ggtctattct	4260
tttgatttat	aagggttttt	gccgatttct	gcctattggt	taaaaaatga	gctgatttaa	4320
caaaaaattta	acgcgaattt	taacaaaata	ttaacgttta	caatttcagg	tggcactttt	4380
cggggaaatg	tgcgcgggaa	ccctatttct	ttatttttct	aaatacatct	aaatatgtat	4440
ccgctcatga	attaattctt	agaaaaactc	atcgagcatc	aaatgaaact	gcaattttat	4500
catatcagga	ttatcaatac	catatttttg	aaaaagccgt	ttctgtaatg	aaggagaaaa	4560
ctcaccgagg	cagttccata	ggatggcaag	atcctggtat	cggctctcga	ttccgactcg	4620
tccaacatca	atacaacctc	ttaattttcc	ctcgtcaaaa	ataaggttat	caagtgagaa	4680

WO 00/41524

PCT/US00/00580

3

atcaccatga	gtgacgactg	aatccggtga	gaatggcaaa	agtttatgca	tttctttcca	4740
gacttggtca	acagggcagc	cattacgctc	gtcatcaaaa	tcactcgcat	caaccaaac	4800
gttattcatt	cgtgattgcg	cctgagcgag	acgaaatacg	cgatcgctgt	taaaaggaca	4860
attacaaaca	ggaatcgaat	gcaaccggcg	caggaacact	gccagcgcat	caacaatatt	4920
ttcacctgaa	tcaggatatt	cttctaatac	ctggaatgct	gttttcccgg	ggatcgagct	4980
ggtgagtaac	catgcatcat	caggagtacg	gataaaatgc	ttgatggtcg	gaagaggcat	5040
aaattccgtc	agccagttta	gtctgacccat	ctcatctgta	acatcattgg	caacgctacc	5100
tttgccatgt	ttcagaaaca	actctggcgc	atcgggcttc	ccatacaatc	gatagattgt	5160
cgcacctgat	tgcccgcacat	tatcgcgagc	ccattttatac	ccatataaat	cagcatccat	5220
gttggaattt	aatcgcggcc	tagagcaaga	cgtttcccgt	tgaatatggc	tcataacacc	5280
ccttgattta	ctgtttatgt	aagcagacag	ttttattggt	catgacccaa	atcccttaac	5340
gtgagttttc	gttccactga	gcgtcagacc	ccgtagaaaa	gatcaaagga	tcttcttgag	5400
atcctttttt	tctgcgcgta	atctgctgct	tgcaaacaaa	aaaaccaccg	ctaccagcgg	5460
tggtttggtt	gccggatcaa	gagctaccaa	ctctttttcc	gaaggtaact	ggcttcagca	5520
gagcgcagat	accaaatact	gtccttctag	tgtagccgta	gttaggccac	cacttcaaga	5580
actctgtagc	accgcctaca	tacctcgctc	tgctaactct	gttaccagtg	gctgctgcca	5640
gtggcgataa	gtcgtgtctt	accgggttgg	actcaagacg	atagttaccg	gataaggcgc	5700
agcggtcggg	ctgaacgggg	ggttcgtgca	cacagcccag	cttggagcga	acgacctaca	5760
ccgaactgag	atacctacag	cgtgagctat	gagaaagcgc	cacgcttccc	gaagggagaa	5820
aggcggacag	gtatccggta	agcggcaggg	tcggaacagg	agagcgcacg	agggagcttc	5880
cagggggaaa	cgcttggtat	ctttatagtc	ctgtcgggtt	tcgccacctc	tgacttgagc	5940
gtcgattttt	gtgatgctcg	tcaggggggc	ggagcctatg	gaaaaacgcc	agcaacgcgg	6000
ccttttttcc	gttccctggc	ttttgctggc	cttttgctca	catgttcttt	cctgcgttat	6060
ccccgtattc	tgtggataac	cgtattaccg	cctttgagtg	agctgatacc	gctcgcgcga	6120
gccgaacgac	cgagcgcagc	gagtcagtga	gcgaggaagc	ggaagagcgc	ctgatcggtt	6180
attttctcct	tacgcacatc	tgccgtattt	cacaccgcac	atatgggtga	ctctcagtag	6240
aatctgctct	gatgccgcat	agttaagcca	gtatacactc	cgtatcgctc	acgtgactgg	6300
gtcatggctg	cgccccgaca	cccgcacaaca	cccgtgacg	cgccctgacg	ggcttgctctg	6360
ctcccggcat	ccgcttacag	acaagctgtg	accgtctccg	ggagctgcat	gtgtcagagg	6420
ttttaccgtg	catcacccga	acgcgcgagg	cagctgcggt	aaagctcatc	agcgtggctg	6480
tgaagcgatt	cacagatgtc	tgccgtgtca	tcgcgctcca	gctcggttag	tttctccaga	6540
agcgttaatg	tctggcttct	gataaagcgg	gccatgttaa	ggcggttttt	ttcctgtttg	6600
gtcactgatg	cctccgtgta	agggggattt	ctgttcatgg	gggtaatgat	accgatgaaa	6660
cgagagagga	tgctcacgat	acgggttact	gatgatgaac	atgcccggtt	actggaacgt	6720
tgtgagggtg	aacaactggc	ggatgggatg	cggcgggacc	agagaaaaat	cactcagggg	6780
caatgccagc	gcttcgttaa	tacagatgta	gggtgtccac	agggtagcca	gcagcatcct	6840
gcatgacaga	tcgggaacat	aatggtgcag	ggcgtgactc	tccgcgtttc	cagactttac	6900
gaaacacgga	aaccgaagac	cattcatggt	gttgctcagg	tcgcagacgt	tttgacgacg	6960
cagtcgcttc	acgttcgctc	gcgtatcggt	gattcattct	gctaaccagt	aaggcaaccc	7020
cgccagccca	gcccgggtct	caacgacagg	agcacgatca	tcgcaccccg	tggggccgcc	7080
atgccggcga	taatggcctg	cttctcgccg	aaacgtttgg	tggcgggacc	agtacgaag	7140
gcttgagcga	ggcggtgcaa	gattccgaat	accgcaagcg	acaggccgat	catcgctcgc	7200
ctccagcgaa	agcgggtcct	gcccgaatat	accagagcgc	ctgccggcac	ctgtcctacg	7260
agttgcatga	taaagaagac	agtcataagt	gcggcgacga	tagtcatgcc	ccgcgccac	7320
cggaaggagc	tgactgggtt	gaaggctctc	aagggcacgc	gtcgagatcc	cgggtgcctaa	7380
tgagtgagct	aacttacatt	aattgcgttg	cgctcactgc	ccgctttcca	gtcgggaaac	7440
ctgtcgtgcc	agctgcatta	atgaatcggc	caacgcgcgc	ggagaggcgc	tttgctgatt	7500
gggcgcacag	gtgggttttc	ttttaccag	tgagacgggc	aacagctgat	tgcccttcac	7560
cgcttgcccc	tgagagagtt	gcagcaagcg	gtccacgctg	gtttgcccc	gcaggcgaaa	7620
atcctgtttg	atggtgggta	acggcggtat	ataacatgag	ctgtcttcgg	tatcgctgta	7680
tcccactacc	gagatatccg	caccaacgcg	cagcccggac	tcggtaatgg	cgcgcattgc	7740
gcccagcgcc	atctgatcgt	tggaaccag	catcgcagtg	ggaacgatgc	cctcattcag	7800
catttgcatg	gtttgttgaa	aaccggacat	ggcactccag	tcgccttccc	gttccgctat	7860
cggctgaatt	tgattgcgag	tgagatattt	atgccagcca	gccagacgca	gacgcgccga	7920
gacagaactt	aatgggcccc	ctaacagcgc	gattttctgg	tgacccaatg	cgaccagatg	7980
ctccacgccc	agtcgcgtac	cgtcttcacg	ggagaaaata	atactgttga	tgggtgtctg	8040
gtcagagaca	tcaagaaata	acgcgggaac	attagtgcag	gcagcttcca	cagcaatggc	8100



WO 00/41524

PCT/US00/00580

4

```

atcctggtca tccagcggat agttaatgat cagcccactg acgcgttgcg cgagaagatt 8160
gtgcaccgcc gctttacagg cttcgacgcc gcttcgttct accatcgaca ccaccacgct 8220
ggcaccagct tgatcggcgc gagatttaac cgccgcgaca atttgcgacg gcgcgtgcag 8280
ggccagactg gaggtggcaa cgccaatcag caacgactgt ttgcccgcca gttgttggtg 8340
cacgcgggtg ggaatgtaat tcagctccgc catcgccgct tccacttttt cccgcgtttt 8400
cgcagaaaac tggctggcct ggttcaccac gcgggaaaac gtctgataag agacaccggc 8460
atactctgcg acatcgata acgttactgg ttccacattc accaccctga attgactctc 8520
ttccgggccc tatcatgcca taccgcgaaa ggttttgccg cattcgatgg tgtccgggat 8580
ctcgacgctc tcccttatgc gactcctgca ttaggaagca gccagtagt aggttgaggg 8640
cgttgagcac cgccgcgcga aggaatggtg catgcaagga gatggcgccc aacagtcccc 8700
cgccacggg gctgcccacc ataccacgc cgaaacaagc gctcatgagc ccgaagtggc 8760
gagcccgatc ttcccatcg gtgatgtcgg cgatataggc gccagcaacc gcacctgtgg 8820
cgccggtgat gccggccacg atgcgtccgg cgtagaggat cgagatctcg atcccgcgaa 8880
attaatacga ctactatag gggaattgtg agcggataac aattcccctc tagaaataat 8940
tttgtttaac ttttaagaag agatatacat 8970

```

<210> 2  
 <211> 19  
 <212> DNA  
 <213> Bacteriophage

<220>  
 <223> BAC-1

<400> 2

cgcggtacac cgacgtcaa

19

<210> 3  
 <211> 19  
 <212> DNA  
 <213> Bacteriophage

<220>  
 <223> BAC-2

<400> 3

cgcggtacac cgacttaat

19

<210> 4  
 <211> 10  
 <212> DNA  
 <213> Bacteriophage

<220>  
 <223> BAC-3

<400> 4

gtcgggtgtac

10

WO 00/41524

PCT/US00/00580

5

<210> 5  
<211> 23  
<212> DNA  
<213> Bacteriophage

<220>  
<223> T7

&lt;400&gt; 5

taatacgact cactataggg cga

23

<210> 6  
<211> 24  
<212> DNA  
<213> Bacteriophage

<220>  
<223> SP6

&lt;400&gt; 6

catacgattt aggtgacact atag

24